






EU Business @ Biodiversity Platform

UPDATE REPORT 1
19 NOVEMBER 2018







Lead Author

JOHAN LAMMERANT Lead Natural Capital and Biodiversity Expert

M +32 495 587 563

E johan.lammerant@arcadis.com

Contributors

LARS MÜLLER

Policy officer at DG Environment, European Commission

E Lars.MUELLER@ec.europa.eu

JEROME KISIELEWICZ

Managing Consultant, ICF

E Jerome.kisielewciz@icf.com







CONTENTS

READE	RS GUIDE	4
1 OB	JECTIVE	5
2 ME	THODOLOGY	6
2.1	Scope	6
2.2	Assessment principles	6
2.3	Uniform assessment template	9
2.4	Assessed approaches	10
3 INI	TIAL FINDINGS OF THE COMPARATIVE ANALYSIS	11
4 MA	IN OUTCOMES OF THE WORKSHOP OF 18 SEPTEMBER 2018	15
4.1	Objectives, agenda and participants	15
4.2	Important findings for the further assessment work	15
ANNEX	1: ASSESSMENT OF BIODIVERSITY ACCOUNTING APPROACHES	
FOR B	USINESS	18
1.	The Global Biodiversity Score (CDC Biodiversité)	18
2.	Biodiversity Impact Metric (CISL)	22
3.	Biodiversity Indicator for Extractive Companies (UNEP-WCMC)	31
4.	Product Biodiversity Footprint (I Care & Consult, Sayari)	35
5.	Biodiversity Footprint approach (ASN Bank)	40
6.	The Biodiversity Return on Investment Metric (by IUCN)	43
7.	The Agrobiodiversity Index (ABD) by Bioversity International	45
8.	The Biodiversity Footprint Calculator (PLANSUP)	50
9.	The LIFE Impact Index + Positive Scoring (by LIFE INSTITUTE)	52
10.	Bioscope (by Platform BEE, Dutch Ministry)	58
ANNEX	2: ANALYSIS OF ASSESSMENTS	60







READERS GUIDE

This report includes:

- Section 1: Description of context and objectives;
- Section 2: Description of the methodological approach;
- Section 3: First findings of comparative analysis:
- Section 4: Outcomes of dedicated workshop (18 September 2018, Brussels);
- Annex 1: Technical fiches of the assessed 'biodiversity metrics for business' approaches;
- Annex 2: Technical tables with comparative analysis.

This report will be updated periodically to reflect evolutions in the development of biodiversity accounting approaches for businesses and financial institutions. This first update contains the following changes compared to the initial assessment report (published on 5 September 2018):

- The structure of the report has changed:
 - A Reader's guide is added;
 - o The technical assessment fiches linked to each approach have been moved to Annex 1;
 - The technical tables with the comparative analysis of approaches have been moved to Annex 2;
- Two technical fiches (nr 7, nr 8) have been updated;
- The comparative analysis has been updated to reflect these recent developments;
- A section has been added (Section 4) with the main outcomes of the workshop dedicated to this assessment exercise organized in Brussels on 18 September 2018.

DISCLAIMER

This document was prepared as part of the EU B@B Platform. However, it reflects only the authors' views, and the European Commission cannot be held responsible for any use which may be made of the information contained therein.







1 OBJECTIVE

Natural capital accounting is rapidly gaining interest, not only at the level of countries or regions, but also at the level of business and financial institutions (FIs). The traditional unidimensional perspective of financial performance (GDP for nations, financial reports for companies) is being replaced by a more multidimensional perspective covering also natural and social capital performances. There is a growing interest in non-financial reporting amongst public authorities, FIs and the business community. Non-financial is becoming the new financial.

The consequence of this tendency is the need for natural capital metrics and robust methodologies. For some environmental issues, such as GHG-emissions and water consumption, these metrics are relatively straightforward and already familiar to businesses. However, the situation is different for biodiversity as key component of comprehensive natural capital assessments. Generally accepted and applied methodologies on how businesses and FIs can measure and value their impacts and dependencies on ecosystem services (ES) and biodiversity are currently lacking.

Businesses are struggling to identify approaches to measure their biodiversity performance that are on the one hand practical and pragmatic and on the other hand meaningful and relevant. This also applies to financial institutions who are looking for suitable ways to assess biodiversity performance (in terms of impacts and risks) of their portfolio of investments.

Why is it so difficult to develop a generally accepted approach to measure and value the biodiversity performance of a company or investment? Multiple obstacles exist, such as the complexity of biodiversity itself (i.e. multitude of species and habitats, each with their specific sensitivity to different pressure categories) and the often complicated cause/impact relationships between a company's activities and biodiversity receptors, to only name a few.

In the light of the growing attention for natural capital accounting (NCA) and a clear demand for methodologies for including ecosystems and biodiversity into NCA, several attempts are ongoing to develop pragmatic biodiversity metrics for businesses and FIs. Faced with these multiple approaches and based on requests from its members, the EU Business @ Biodiversity Platform (EU B@B Platform)¹ conducted a critical and constructive assessment of a sample of biodiversity accounting approaches developed for or by businesses and FIs. As a first step the aim of this exercise is to provide an overview of different methodological approaches and compare their key features. The second objective is to identify the key obstacles faced by these approaches and the gaps that still need to be filled. Considering the business demand for such methodologies, findings are likely to inform the future activities of the EU B@B Platform and potential partners (including developers and users of these metrics). This work shall also feed into the initiative to prepare a Biodiversity Supplement to the Natural Capital Protocol.

This assessment exercise will be regularly updated. This report is the first of a series of Update Reports which will be prepared on behalf of the EU B@B Platform. Updates might consist in the inclusion of additional approaches in the assessment, adaptations of the assessment of the approaches are already included to reflect new developments, descriptions of case studies, etc. We therefore invite further constructive contributions by other members of in the EU B@B Platform and beyond with a view to continuously improve insights and solutions in the field of biodiversity metrics for the private sector.

Important: We would like to stress that this discussion and updated assessment paper is only an intermediary step and is neither meant to be definitive nor exhaustive. We rather hope to facilitate a discussion based on enhanced transparency with a view to facilitate the further development and enhancement of these much-needed methodological initiatives.

¹ in particular Workstream 1 on Natural Capital Accounting and Workstream 3 on Financial Institutions and Biodiversity



_





2 METHODOLOGY

2.1 Scope

Given the wealth of biodiversity assessment approaches, clear boundaries were set to scope this assessment: the focus of this assessment is on biodiversity accounting approaches for businesses and financial institutions (Fls) which rely on quantitative indicators that provide information on the significance of impacts on biodiversity, and which are not case-specific. The latter is important, as businesses and Fls need approaches that can be applied by several companies and preferably for different types of business applications, different levels of application (e.g. project, site) and in different locations.

For this reason and without any prejudice to their value and usefulness, certain types of biodiversity assessment approaches are not included in this assessment, such as:

- Process based approaches which rely on 'yes/no' questions and only provide qualitative insights on
 the level of actions undertaken by a company in the field of biodiversity. They rely on 'process
 indicators' (e.g. 'Do you have a biodiversity action plan?') rather than 'impact indicators'. Examples
 of such approaches include the European Biodiversity Standard², the Biodiversity Benchmark³, and
 the Biodiversity Check⁴;
- Approaches applied in Environmental Impact Assessment and similar types of specialized studies, which focus on a specific development in a specific area.

Other approaches that were not included to date in this exercise include biodiversity metrics approaches which can be used for calculating biodiversity offsets. Examples are the BBOP biodiversity metrics⁵, the UK DEFRA biodiversity metrics⁶, the Dutch 'Natuurpunten' method⁷, etc. These latter approaches could also fit under the given scope but were not included to date.

2.2 Assessment principles

The key principle underpinning this assessment is the applicability of the biodiversity metrics and related tools for businesses and FIs. Therefore, it's important to link the approaches on biodiversity metrics to the **type of applications** businesses and FIs are interested in as well as to **the scope** (e.g. product level or site level) (see Table 1). Table 1 presents some examples of such 'business applications'. An example of how a business application can be different according to its scope (level of application) is 'comparing different options':

- A financial institution looking for ways to make its investment portfolio more sustainable, might be interested in the biodiversity performance of different sectors or companies (corporate level)
- A multinational company might be interested in the biodiversity performance of each site, or in the biodiversity performance of its products
- A company site manager might be interested in comparing two different projects (e.g. locations, technologies)

⁷ https://www.pbl.nl/publicaties/2009/natuureffecten-in-de-mkba-s-van-projecten-voor-integrale-gebiedsontwikkeling



² http://www.europeanbiodiversitystandard.eu/en

³ https://www.wildlifetrusts.org/sites/default/files/2018-06/BBO M4%20Biodiversity%20Benchmark%20Requirements.pdf

⁴ https://www.business-biodiversity.eu/docs/ebbc_index01.aspx?id=36799&basehrefregu=true&isalias=true

⁵ http://bbop.forest-trends.org/pages/guidelines

⁶ https://www.gov.uk/government/publications/technical-paper-the-metric-for-the-biodiversity-offsetting-pilot-in-england





Table 1: Rusiness	annlications	where	'hiodiversity	nerformance'	is an	important indicator

Business application	Product level	Project level	Site level	Corporate	Supply chain level	Investment portfolio
Assessing biodiversity risks and/or opportunities Assessing nature and magnitude of biodiversity impacts and dependencies, and their associated business risks and opportunities	Х	Х	Х	Х	Х	
Comparing options Compare, contrast and select from a range of alternative options, while considering their relative biodiversity performance	Х	Х	Х		Х	X
Going for No Net Loss or Biodiversity Net Gain Assessing net biodiversity impact; this requires a sound understanding of the baseline situation	Х	X	Х	Х	X	X
Communicating internally or externally Communication on biodiversity performance, and if relevant evolution over time	X	Х	Х	Х	Χ	Х

Other?

Source: NCP, CoP Finance @Biodiversity of the of EU B @B Platform8.

Furthermore, the assessment approach is underpinned by the same principles as applied in the NCP, i.e.:

- 1. Relevance:
- 2. Rigor;
- 3. Replicability; and
- 4. Consistency.

Relevance implies that the most relevant biodiversity issues are covered, i.e. those that are most material for the business or the FIs and their stakeholders. This requires:

- A clear insight in the cause/impact relationships, i.e. which pressures are causing which impacts on which biodiversity receptors? The required materiality⁹ and level of detail need to be adequate to each specific situation, and this is often determined by stakeholder expectations.
- A correct set of indicators, which provide relevant information to inform business decisions; indicators need to be responsive to changes (i.e. pressure indicators need to reflect changes in pressures, while species or habitat indicators should be able to reflect changes in the state of biodiversity as a result of company actions).

⁹ an impact or dependency on natural capital is material if consideration of its value, as part of the set of information used for decision making, has the potential to alter that decision (Natural Capital Protocol; Adapted from OECD 2015 and IIRC 2013).



⁸ During the workshop of 21 March 2018, the CoP Finance (Workstream 3 of the EUB@B Platform) proposed the following specific types of business applications for Fls:

portfolio level assessments (Fls need KPls on biodiversity impacts/performance by sector)

company level assessments (Fls often apply 'best in class' approaches for benchmark purposes, so there is a need for reference benchmarks)

project level assessments

internal purposes vs reporting purposes





A key challenge in this respect is the quest for a simple and comprehensive biodiversity indicator. Does it really exist? Is such indicator compatible with the abovementioned criterion of relevance? A complicating factor is the duality of the term 'biodiversity', as it comprises both intrinsic biodiversity values (i.e. species, habitats, and their functional relationships) and ES. Businesses can have impacts on ES and might also be dependent on ES. This assessment considers both elements of biodiversity.

Several approaches rely on one overall proxy indicator for biodiversity. In such case, the type of applications will be limited.

Rigor refers to the use of technically robust (i.e. from a scientific and economic perspective) information, data and methods that are also fit for purpose. This requires:

- The use of up-to-date information on the presence and sensitivity of species and habitats within the influence area of company activities. The required level of details is often determined by stakeholder expectations.
- Correct information on the baseline situation e.g. in case of project appraisal.

Replicability means that all assumptions, data, caveats, and methods used are transparent, traceable, fully documented, and repeatable. This requires that approaches don't contain black boxes.

Consistency is the principle that all data and methods used for an assessment are compatible with each other and within the scope of the analysis, which depend on the overall objective and expected application (e.g. option appraisal, assessing biodiversity risks and opportunities, aiming for no net loss or biodiversity net gain).

Additional assessment parameters, which are relevant in the context of this assessment exercise, include10:

- Sector coverage: Is it sector specific or can it be used by several sectors?
- User friendliness:
 - o Is it easy to use by non-experts or does it require specialist knowledge?
 - O What are the required time efforts?
 - o Does it require a similar approach as other environmental issues or is it very specific?
- Sources of information feeding into the approach (i.e. is it based on real data (based on observations or measurements) or modelling?)
 - For model-based outcomes, it is very useful to know if real data (on pressures, on state of biodiversity) can be plugged in, in order to refine these outcomes:

 $^{^{10}}$ Apart from 'sector coverage', the other additional criteria were suggested during the CoP Finance (WS 3) workshop of 21 March 2018 in Brussels







2.3 Uniform assessment template

To review the approaches and initiatives selected for this assessment in a uniform and objective way their developers were invited to fill in a common template and provide answers to the following questions and elements:

- Title/Name of initiative:
- Date of assessment, including updates;
- Description of involved actors, and type of process;
- Description of state-of-the-art of current developments (starting development of methodology, pilot phase, already (widely) applied in sector, etc.);
- Description of the metric(s) and methodology behind it;
- Outcomes informed by real data or by modelling?
- Description of types of business application (see Box 1), i.e. what kind of decisions could be based on it? The following applications are considered:
 - At which levels of the value chain can the approach be used: site level, corporate level, supply chain level, product level?
 - Can the approach be applied to compare different investment options?
 - Can the approach be used to assess progress of a biodiversity action plan?
 - o Can the approach be used to proceed towards no net loss or biodiversity net gain?
 - o Does the approach only apply to new developments or does it also cover existing activities?
 - Etc...
- Which sectors are covered by current initiatives?
- Which drivers of biodiversity loss are covered e.g. land use;
- Level of detail of data, data sources, flexibility in granularity, etc.;
- User friendliness:
- Strengths and weaknesses, according to initiators.

The completed templates are included as technical fiches under Annex 1. These are numbered according to the sequence in which they have been assessed.







2.4 Assessed approaches

Within the scope of this assessment, several initiatives have been identified based on research and bilateral consultations with developers of assessed approaches (who often started the development of their approach with an inventory of existing approaches). Most of these approaches are still under development. Table 2 provides an overview. A detailed description of each approach is included in Annex 1.

Table 2: Overview of biodiversity accounting approaches included in this assessment report

	Developer	Name of tool	Assessment
1	CDC Biodiversité	Global Biodiversity Score	Completed and assessed by EU B@B Platform in Jan 2018, including replies from CDC Final update on 22 August 2018
2	Cambridge Institute for Sustainable Leadership (CISL)	Biodiversity Impact Metric	Completed on 15 March 2018 Reviewed by EU B@B Platform Update on 20 August 2018
3	UNEP-WCMC	Biodiversity Indicators for Extractive Companies	Completed on 19 April 2018 Reviewed by EU B@B Platform Update on 15 July and 21 August 2018
4	I CARE - Sayari	Product Biodiversity Footprint	Completed on 20 June 2018 Reviewed by EU B@B Platform. Update on 23 August 2018
5	ASN Bank	Biodiversity Footprint approach	Completed on 28 June 2018 Reviewed by EU B@B Platform Update on 24 August 2018
6	Bioversity International	Agrobiodiversity Index (ABD)	Completed on 1 August 2018 Reviewed by EU B@B Platform
7	Plans Up	Biodiversity Footprint Calculator	Information from website
8	LIFE Institute	Impact Index + Positive Scoring	Completed on 21 August 2018 Reviewed by EU B@B Platform Update on 2 November 2018
9	Platform BEE (Dutch Ministry)	Bioscope	Information from website Update by Plansup on 4 November 2018
10	IUCN	Biodiversity Return on Investment Metric (BRIM)	Completed on 29 August 2018 Reviewed by EU B@B Platform

Source: EU B@B Platform, 2018.

From this list, nine assessment forms were completed by the developers themselves. These assessment forms were then adapted and completed based on a critical review completed by the NCA Workstream Leader of the EU B@B Platform. Approach nr 9 was based on publicly available information, as time was lacking to get in touch with the developers. The information provided for this approach should therefore be considered with caution.

We are very grateful for the very constructive cooperation with all contacted developers.







3 INITIAL FINDINGS OF THE COMPARATIVE ANALYSIS

The comparative analysis of the biodiversity accounting approaches for businesses and FIs selected for this assessment revealed the following findings:

- Most approaches are under development and (will) make use of a road-testing phase.
- Results from road-testing are currently scarce. Although some approaches are operational no case studies have been published to date. As the 'proof of the pudding is in the eating' it will be crucial to integrate case study findings in the next phase of this assessment exercise.
- Approaches are not only being developed for the business community but also for the finance community, which is a very positive development and reflects the growing awareness that FIs can play a key role in pushing company performance on biodiversity upward.
- Most approaches are applicable across sectors. The only exceptions to this rule are the UNEP-WCMC approach¹¹ that only addresses the extractives sector; the ABD Index¹² which is only applicable to the agri-food sector (and FIs investing in it); and the CISL approach¹³ which focuses on land use in the supply chain.
- Some developers have clearly articulated the key principles for suitable biodiversity metrics for businesses. As an example, the key principles developed by CISL are listed in Table 3, but CDC Biodiversité¹⁴ and UNEP-WCMC have also developed a set of principles. Despite some differences they are largely overlapping.

Table 3: Principles for a suitable biodiversity metric (Source: CISL)

Principle	Description
Meaningful	Meaningful to business and investor communities so it can be used to drive decision making. Methodology is clearly understood.
Measurable and comparable	Allows for comparison across geographies and time.
Possible to aggregate	Can be aggregated from site-level to regional and global scales.
Practical	Data is accessible, measurable by company or using free, globally available data. Ability to substitute better information where available.
Replicable and credible	Based on a reputable scientific method.
Context based	Considers local conditions/levels to reflect 'impact' (beyond 'usage').
Responsive	Responds to changes in company activities, both short and long term.

- Biodiversity accounting approaches generally follow the same logic for identifying impacts:
 - Step 1: scoping (economic activities, products);
 - Step 2: linking economic activities to pressures;
 - Step 3: linking pressures to biodiversity impacts. This requires coefficients for linking pressures to impacts, as well as data on biodiversity in the affected areas.

Steps 2 and 3 often rely on the same underpinning models or data sources, i.e.:

For Step 2: the Exiobase matrix-based input-output model¹⁵ (economic activities → pressures);

12 Fiche nr 6

¹⁵ http://www.exiobase.eu/



11

¹¹ Fiche nr 3

¹³ Fiche nr 2

¹⁴ Fiche nr 1





 For Step 3: the GLOBIO¹⁶ (global estimates of biodiversity abundance) and ReCiPe¹⁷ (Life-Cycle Assessment) for linking pressures to impacts; the GLOBIO, IUCN Red Lists¹⁸ and IBAT¹⁹ for assessing biodiversity values

As a result, the strengths and weaknesses of the developed approaches are to a large extent defined by the intrinsic strengths and weaknesses of these models and data sources.

Some examples of remaining weaknesses related to GLOBIO include (ReCiPe has similar weaknesses):

- Pressure-impact relationships in the GLOBIO model are biased towards the most studied species and ecosystems;
- Marine biodiversity is not factored in;
- Overexploitation, invasive species, chemical pollution and soil degradation are not factored in yet;
- GLOBIO is not species and habitats specific.

Other shortcomings were flagged with regard to linking pressures to economic activities. One of the potential shortcomings of this step is the overemphasis on the IUCN Red List species, Critical Habitats and 'protected areas', which might mask the ongoing degradation of more common biodiversity. Therefore, wherever available additional data should be included in the accounting approach (e.g. local Red Lists, indicator species representing common biodiversity).

- Some approaches acknowledge the need to introduce real life data into their methodologies, either
 by replacing modelled data or by completing modelled data with real life data (hybrid approaches). A
 critical attitude at the side of the developers is key. Metrics showing nice biodiversity
 performance, but relying on assumptions which don't reflect reality, or which are based on inaccurate
 descriptions of measures (e.g. moving from minimal to light and intensive land use management) are
 to be avoided.
- Most approaches foresee the possibility to adapt score performance according to the actions that are taken (i.e. biodiversity actions are directly reflected in the scores). In those cases, businesses should make sure to align the type of decisions they are taking with the granularity level of the input data (i.e. economic activities, pressures, biodiversity values). Strategic decisions might only require high-level information that provides insights if key components of biodiversity will improve or degrade and what the company can do to change this (e.g. selection of new areas for commodity production). Other decisions might require more detailed information even at species level, for instance in the case of reputational risks.
- Most approaches rely on one single biodiversity metric, with MSA (Mean Species Abundance) and PDF (Potentially Disappeared Fraction of species) being the most popular ones. I Care and Sayari's PBF approach²⁰ uses PDF but for each impact driver separately (spider diagram). IUCN's BRIM approach²¹ relies on one single metric too, but the scope of this tool is very specific. The UNEP-WCMC approach for extractive industries aims for a 'set of site level indicators, allowing aggregation at corporate level (to be developed). It is interesting to note that the working group accompanying this initiative decided that a single metric may not provide the necessary resolution, flexibility or information to meet the indicator needs of the extractive sector. The ABD Index is based on 33 indicators.
- All approaches score relatively well in terms of rigor, replicability and consistency. The main challenge seems to be linked to 'relevance' criteria. Relevance means that the most relevant

²¹ Fiche nr 10



¹⁶ http://www.globio.info/

¹⁷ https://www.pre-sustainability.com/recipe

¹⁸ https://www.iucnredlist.org/

¹⁹ https://www.ibatforbusiness.org/

²⁰ Fiche nr 4





biodiversity issues are covered, i.e. those that are most material for the business and its stakeholders (Source: NCP). As mentioned under section 2.2 this requires a clear insight in the cause/impact relationships (i.e. which pressures are causing which impacts on which biodiversity receptors?) and a correct set of indicators, which provide relevant information to inform business decisions e.g. indicators need to be responsive to changes. As most metrics are based on models (see above), they tend to miss the local context of each specific case. Moreover, as and are comprehensive (see above)

- **ES** are generally not included in the accounting approaches (GBS assumes high ES values when intrinsic biodiversity values are high), apart from the approach by LIFE Institute²², which considers impacts on ES too.
- In terms of **coverage of business applications throughout the value chain**, all parts of the value chain are covered by one or more of the assessed approaches, which is a positive observation:
 - o 4 approaches cover a product level assessment;
 - Only 2 approaches cover project level assessments, one of them limited to large scale projects;
 - 3 approaches cover site level assessments;
 - 4 approaches cover the supply chain;
 - 6 approaches cover the corporate level;
 - o 7 approaches might be applicable by FIs.

Some approaches (Global Biodiversity Score, ABD Index) are also suitable for country level assessments, which is out of scope for this study, but these offer promising potential in the light of the increased demand for better alignment between business and national level NCA.

None of the approaches covers all types of business applications.

- As presented in Table 4, six different types of business applications are covered by one or more biodiversity accounting approaches. 'Assessing biodiversity performance' always allows 'comparing options', so the numbers for these two applications are equal. This also applies to 'communication'. 'No Net Loss' applications and 'offering financial products based on high biodiversity performance' are interesting additional business applications, which are only covered by a minority of approaches.
- Most methods are not particularly aligned with other NCA approaches. The Global Biodiversity Score (GBS) of CDC Biodiversité does aligned with metrics used in the GHG Protocol²³ as it acknowledges the importance of more convergence between biodiversity accounting approaches and NCA approaches. Carbon footprinting data and results can be re-used as direct inputs for the GBS and the approach is very similar, both in terms of data collection and of concepts (e.g. scopes 1, 2 and 3, attribution of responsibilities across the value chain). The Healthy Ecosystem Metric of CISL aims to integrate biodiversity, soil and water performances of land use activities into one overall metric. The Natural Capital Impact Matrix of the LIFE Institute includes waste generation, GHG emission, water consumption, energy use and the area occupied by the business.

²³ https://ghgprotocol.org/



²² Fiche nr 8





Table 4: Number of approaches covering different types of business applications

Business application	Product level	Project level	Site level	Corporate	Supply chain level	Investment portfolio
Assessing biodiversity performance (risks and/or opportunities) Assessing nature and magnitude of biodiversity impacts and dependencies, and their associated business risks and opportunities	4	2	3	6	4	6
Comparing options Compare, contrast and select from a range of alternative options, while considering their relative biodiversity performance	4	2	3	6	4	6
Going for No Net Loss or Biodiversity Net Gain Assessing net biodiversity impact; this requires a sound understanding of the baseline situation	1	1	1	2	0	2
Communicating internally or externally Communication on biodiversity performance, and if relevant evolution over time	4	2	3	6	4	6
Assessing Rol of investments in biodiversity restoration	0	0	1	1	0	1
Offering financial products based on high biodiversity performance (e.g. green bonds)	0	0	0	0	0	1







4 MAIN OUTCOMES OF THE WORKSHOP OF 18 SEPTEMBER 2018

4.1 Objectives, agenda and participants

On 18 September 2018 a dedicated workshop "Biodiversity Metrics for Business and Finance" was organized by two Workstreams of the EU Business and Biodiversity Platform: CoP
Finance@Biodiversity and the Natural Capital Accounting workstream, and was open to developers and practitioners in the emerging area of biodiversity accounting.

The workshop's specific objectives were to:

- Inform on available and emerging 'biodiversity metrics for business and finance' approaches, based on analysis of pro's and con's, and gain deeper understanding of what is considered within each metric;
- Explore applicability of metrics by practitioners. 'To what extent are the current metrics under development meeting the needs of business and finance?';
- Identify complementarities between approaches, as well as remaining gaps and obstacles, and potential ways to deal with them;
- Inform related needs for further work over the next few years, as well as future research and innovation needs (e.g. the upcoming Horizon2020 project on NCA for business).

More details about the workshop including the minutes and presentations can be found on the website of the EU B@B Platform²⁴.

4.2 Important findings for the further assessment work

Although all workshop findings are useful and interesting on their own, a number of them are of particular importance for the further assessment exercise. The below overview in Table 5 is a selection of the outcomes of the workshop. Some of these findings are useful observations which could be included in Section 3 of this report, while other findings might be inspirational for further action, either for future research (by the Platform, by developers, supported by funding e.g. H2020) or for strengthening existing cooperation or establishing new cooperation (EU B@B Platform with other stakeholders).

Table 5: Key findings of the workshop and relevance for the metrics assessment work

Main findings	Relevance
Most methodologies are developed via collaboration between academics and business (including finance)	Info for report
Sector specific approaches are included in the assessment report, but it's worth looking at company specific approaches too	Future work Platform?
Most approaches focus on (avoiding and minimizing) negative impacts. Dependencies and positive impacts are still too much under the radar.	Research need
Ideas for additional approaches to be explored: • framework that could be used to valuate positive impact, partly based on TEEB, (Wageningen University) • AERM project which combines LCA and ESS valuation; it provides a link to a database. It includes ESS dependency between stocks, materials, and flows (see http://www.naturalcapitalfinancealliance.org/aerm-project)	Future update report
Many approaches are high level and serve as a compass to identify biodiversity hotspots (in the sense of focus areas for the company to work on). A stepwise approach from 'Shine a light' to 'Deep dive' and 'Focus action' is preferred. Need for maximum complementarity	Research on complementarities between approaches

²⁴ http://ec.europa.eu/environment/biodiversity/business/news-and-events/news/news-63 en.htm



15





between high level scale and local scale and data. Need for better alignment macro- and micro scale approaches (tiered approach)

In terms of the link between natural capital and biodiversity, first focus on natural capital Research need and then to zoom in on biodiversity Apart from the business applications that were identified in the report, workshop participants came up with further specifications and additional applications: ESG screening of equity / investments (more specific type of application for FI, under 'Assessing biodiversity performance' business application in report (Table 4: Number of approaches covering different types of business applications) The scope 'product level' needs to comprise services as well Shared resources or ES with other stakeholders (impact on choice of metrics?) Communication and reporting Biodiversity footprinting of products as input for setting a price (or tax); this is an Info for report interesting business application, similar to 'Offering financial products based on high biodiversity performance' (see Table 4: Number of approaches covering different types of business applications), but now on product level Compliance to legal obligations or FI safeguards Providing more transparency (to be added to 'Communicating internally and externally' (Table 4: Number of approaches covering different types of business applications) To feed into the Sustainable Finance Taxonomy (EU process) Recently University of Oxford in cooperation with IUCN has elaborated a taxonomy on Need for alignment and business applications. It's worth exploring consistency with this taxonomy. cooperation The Biodiversity Footprint collaboration of EU CoP Finance@Biodiversity members ASN Bank, Actiam, CDC Biodiversité and Finance in Motion is shaping up under the name of Future update report 'Biodiversity Footprint Financial Institutions' (BFFI); A remaining challenge for the BFFI group is how to include positive impacts; for example by including parameters such as certifications in the rating of biodiversity performance of Research need sectors/companies e.g. FSC certification, RSPO (Round Table Sustainable Palm Oil); Based on the footprint results, engagement with investees might be more powerful than shifting investment to another investee (exclusion)! Shifting investment to less damaging Info for report sectors or companies, doesn't solve the biodiversity impact problem which those investees cause. Assessment work by EU Soon, the 'Common Ground for Biodiversity Accounting' joint paper by the four EU CoP B@B Platform used as F@B members (ASN Bank, Actiam, CDC Biodiversité, Finance in Motion) will be one of the underpinning published. ASN Bank will include their whole life support system thinking, connecting sources: biodiversity to human rights, the SDGs and the Planetary Boundaries concept; Info for report: Continued cooperation Research on The PBF approach (spatialized LCA) has complimentary elements to offer to ReCiPe complementarities between approaches Business and finance are looking for a science based target for biodiversity (such as 1,5 C temp increase for climate change). Challenge is to define the reference, and to know what is 'enough' by means of biodiversity impact. The narrative should be about 'aiming for Research need positive impact' and agreed internationally as with the 'Paris agreement for CC'. Positive Need for cooperation impact needs to be more articulated. Planetary Boundaries, the CBD CoP 2020 in Beijing and on-going work by the World Resource Institute could offer opportunities for joint next



Efforts on biodiversity metrics should be linked to the NCC work on the Biodiversity

steps.

Supplement

Continued cooperation





Data quality (lack of standardisation/baseline, location-specificity of biodiversity, company specific data) and availability/price of sufficient data need to be solved for improving biodiversity footprinting of investments, projects and products. Next to transparency on uncertainties, business, finance, biodiversity experts and NGOs should collaborate to provide more and better data for decision-making.

Research need

- There was broad consensus on the following:
 - Need for better alignment business tools and FI tools
 - Need for tools that allow integration of real data
 - Complementarities between tools need to be better articulated
 - Need for guidance on which tools to choose for which company needs (business applications)
 - Need for guidance on how to connect to other NCA approaches (e.g. carbon footprinting, water footprinting) or tools that are aligned with other NCA approaches
 - Need for tools that connect to global biodiversity objectives such as the SDGs or Planetary Boundaries in order to set and achieve more science-based targets
 - Need for guidance on how to integrate dependencies, positive impact and ES, and as such connect to social issues as well (e.g. health)
 - Need for more and more detailed data
 - Need for alignment between data requirements at business level and at national level.

Research need and future work by the Platform







ANNEX 1: ASSESSMENT OF BIODIVERSITY ACCOUNTING APPROACHES FOR BUSINESS

1. The Global Biodiversity Score (CDC Biodiversité)

FICHE NR 1	GLOBAL BIODIVERSITY SCORE (GBS)
Date of assessment	 First assessment submitted to CDC Biodiversité on 12 January 2018 Replies by CDC Biodiversité on 24 January 2018, which were all integrated in the text below
	Update by CDC Biodiversité on 22 August 2018
Actors	<u>Lead</u> : CDC Biodiversité (France) <u>Other</u> : Club of Businesses for Positive Biodiversity (B4B+ Club) acts as a platform for the GBS development (a group of +30 businesses representing different sectors, including finance sector)
Process and current position	CDC Biodiversité launched the biodiversity equivalent of the Teq CO ₂ for climate change in partnership with businesses and financial institutions. It is based on internationally recognised scientific research.
	An excellent description of the GBS approach and state of the art can be found under http://www.mission-economie-biodiversite.com/wp-content/uploads/2017/11/N11-TRAVAUX-DU-CLUB-B4B-INDICATE UR-GBS-UK-BD.pdf . The GBS will remain under development until early 2020: the first biodiversity footprint audits will be conducted in 2020. • Technical developments are ongoing: aquatic drivers impacting biodiversity are being included, the scope of raw materials analysed is expanded and the link between company's activities and raw materials is refined. • The operational relevance of the footprint is being tested thanks to the involvement of future users, i.e. businesses. This happens both through meetings of the B4B+ Club and through several ongoing (and future) case studies, including on sourcing comparison for agricultural commodities and on (investment and loan) portfolio impact assessment
Key features of methodology	GBS™ is designed to provide an overall and synthetic vision of the biodiversity footprint of economic activities. It is not intended to replace local indicators which are best suited to local or on-site biodiversity assessments. This idea of reconciling different scales is key and it is essential that the GBS™ results are consistent with analyses conducted on a local scale, making it possible to summarize the data while losing as little information as possible. The GBS business biodiversity footprint can be estimated in a two-step process (see figure below). First, pressures caused by specific economic activities on biodiversity have to be quantitatively assessed. To analyse the value chain, the GBS™ methodology mainly uses the Exiobase matrix-based input-output model and direct data on pressures when available. Then, the impacts of these pressures on ecosystems have to be estimated. This last step relies on the GLOBIO model which is based on pressure-impact relationships.
	The GLOBIO model, developed by a consortium formed in 2003 consisting of PBL, UNEP GRID-Arendal and UNEP-WCMC, calculates the impact of environmental drivers on biodiversity in the past, present and future. It draws on driver-impact links found in scientific research (pressure – impact relationships are derived from peer-reviewed literature (nearly 300 articles) using meta-analyses). GLOBIO provides both pressure-impact relationships and global estimates of the abundance of biodiversity, based on spatialized data on various environmental drivers (see below) – and not field data on species – as input data. These drivers are taken mainly from the Integrated Model to Assess the Global Environment (IMAGE). GLOBIO produces spatialized results for land







and aquatic (freshwater) biodiversity at a resolution of 0.5° by 0.5°, i.e., 50 km by 50 km at the Equator. These are expressed in terms of average abundance of a species (i.e. MSA). **PRESSURES IMPACTS** LIFE CYCLE ANALYSIS • I and-ueo · Encroachment and fragmentation INPUT-OUTPUT MODELS GLOBIO MODEL Climate change **Biodiversity** Pollutions Invasive exotic species NVESTMENT Resources overexploitation Global **B**iodiversity core Company's and financial actor's biodiversity footprint in km2MSA Rating Metrics High level proxy indicator (at macro-economic level) Principles: Quantitative ('a number') Entire value chain coverage Concise ('which is good for communication purposes') Measures intrinsic biodiversity value (not ES, but is assumed that ES flourish when intrinsic values are OK) Consensual (developed with and accepted by scientists, NGOs, businesses) Complement and be compatible with local indicators GBS uses mean species abundance (MSA) and its surface area equivalent, i.e., km² MSA. The latter is the product of MSA multiplied by the area to which it applies (expressed in km²). MSA measures biodiversity intactness relative to its abundance in undisturbed ecosystems. A 100% ratio indicates an intact ecosystem while damages caused by an increase of pressures bring the MSA progressively to 0% when all originally occurring species are extinct in the ecosystem. Outcomes Hybrid. Modelled results of GLOBIO are used to estimate average industry pressures based on (and impacts) when real data are not available (= 'default assessment'). When real data modelling or on pressures are available, they are instead combined to the pressure-impact real data? relationships provided by GLOBIO to conduct a 'refined assessment'. At each stage, the best available data are used, with a preference for real data over modelled average values. **Drivers** of Drivers (GLOBIO) include: land conversion, fragmentation, encroachment, atmospheric N biodiversity deposition (eutrophication) and climate change for terrestrial biodiversity, and wetlands loss conversion, local and network land-use in catchment of wetlands, hydrological disturbance of wetlands and rivers, land-use in catchment of rivers and eutrophication of lakes for aquatic biodiversity. Missing drivers will be added to GLOBIO through future developments. **Application** The GBS is suitable for the following applications: Calculating the footprint of a financial asset portfolio as a basis for portfolio investment decisions by finance sector, as long as it remains at the level of sectors or companies (when enough company-specific data is available) Corporate level assessments, including estimation of biodiversity performance along the whole value chain of a company. The GBS can







	also be used by non-financial companies themselves to assess a refined
	footprint if detailed information is available. • Country level assessments (out of interest for business biodiversity
	performance assessments)
	 The GBS is not the most suitable for the following applications: Assessments at site level and project level
	Due to:
	 The uncertainties associated to the pressure-impact statistical relationships, especially at small scales (under 1000 ha). Project level assessments are robust only for very large-scale projects (impacting thousands of hectares) and should generally be limited to internal purposes
	 Not species or habitat specific The GBS metrics can only be used for high level indications but at this
	 moment cannot be disaggregated to e.g. site or project level. The GBS could be used for No Net Loss approaches at the corporate or financial institution-level:
	 As any assessment tool, it would require as an input a baseline scenario to compare against the biodiversity losses and gains.
	 Since the GBS is not the most suitable at site or project level and since it focuses on intactness and not on individual species or habitats, it is not appropriate to conduct regulatory No Net Loss project assessments (which usually require particular attention for endangered and protected
	species). o It could however provide balance expressed as net gains or losses of km² MSA if fed with appropriate data.
	applies both to new and existing operations
	assessments can be used for reporting purposes
Required data	The GBS can work with different datasets listed below, by increasing order of usefulness in terms of the precision that can be expected from assessments:
	Economic activity data: turnover by country and industry (of the company consequence of the company of inequal inequal inequal in the company
	assessed or of the company a financial institution invested in);
	 assessed or of the company a financial institution invested in); Pressure data: Carbon emissions on scope 1, 2 and 3
	 assessed or of the company a financial institution invested in); Pressure data: Carbon emissions on scope 1, 2 and 3 Land use changes (ideally using a 13 habitat types nomenclature including
	 assessed or of the company a financial institution invested in); Pressure data: Carbon emissions on scope 1, 2 and 3 Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) Comprehensive biodiversity direct data: when very detailed ecological
	 assessed or of the company a financial institution invested in); Pressure data: Carbon emissions on scope 1, 2 and 3 Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly
	 assessed or of the company a financial institution invested in); Pressure data: Carbon emissions on scope 1, 2 and 3 Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) Comprehensive biodiversity direct data: when very detailed ecological
Granularity	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes:
Granularity level	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio
	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country
	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country For internal purposes:
	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country For internal purposes: - Project or site (indicative values with uncertainties)
level	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country For internal purposes: - Project or site (indicative values with uncertainties) The pressure-impact relationships become imprecise below areas of about 1000 ha.
	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country For internal purposes: - Project or site (indicative values with uncertainties)
level	assessed or of the company a financial institution invested in); • Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) • Comprehensive biodiversity direct data: when very detailed ecological monitoring data are available, the mean species abundance might be directly calculated. For reporting purposes: - Portfolio - Corporate entity or business unit - Country For internal purposes: - Project or site (indicative values with uncertainties) The pressure-impact relationships become imprecise below areas of about 1000 ha. • Is it easy to use by non-experts or does it require specialist knowledge? Using the results of the GBS does not require specialist knowledge and the metrics of km² MSA is relatively easy to understand and visualize (the total land area of the Earth is 130 million km² and about 40 million km² MSA have been lost, i.e. the combined area of Africa and Europe, which is something anyone can easily understand).







	The level of efforts depends on the desired level of details, as the GBS can be flexible: quick approximations can be obtained with industry and country-level averages, and more refined assessments can be obtained if more precise data are collected. A quick assessment takes a couple of weeks and uses easily accessible (and existing) data. A typical detailed assessment should require a couple of months and might require the aggregation (or creation) of additional data, e.g. on habitat maps.
	 Does it require a similar approach as other environmental issues or is it very specific?
	Carbon footprinting data and results can be re-used as direct inputs of the GBS and the approach is very similar, both in terms of data collection and of concepts (e.g. scopes 1, 2 and 3, attribution of responsibilities across the value chain).
Strengths recognised by initiators	 Scientifically well underpinned (best available knowledge and tools e.g. GLOBIO, Exiobase) Quantitative (and scientifically robust) link between pressures and impacts Covers all types of biodiversity and thus adequately captures the trends and risks faced by biodiversity and associated ecological functionality Spatially explicit Covers most drivers for biodiversity loss Covers all industry sectors and all countries Compatible with site-level data (micro) and international objectives (macro) Biodiversity input data (MSA, pressure – impact relationships) based on extensive meta-analysis which continuously allows for adding new studies Allows for introducing weight factors differentiating ecosystem condition based on protection regime, protected species, etc.
Weaknesses recognised by initiators	 Pressure-impact relationships in the GLOBIO model are biased towards the most studied species and ecosystems. Marine biodiversity is not factored in Overexploitation, invasive species, chemical pollution and soil degradation are not factored in yet Remaining shortcomings in reallocation rules (i.e. linking pressures to economic activities)







2. Biodiversity Impact Metric (CISL)

FICHE NR 2	Biodiversity Impact Metric
Date of	First assessment by CISL on 15 th March
assessment	Update on 20 th August
Actors	Lead: Cambridge Institute for Sustainability Leadership (CISL)
	Members of the Natural Capital Impact Group:
	Kering, ASDA, Mondi, Volac; Mars; The Crown Estate; Anglian Water; Yorkshire water; Primark
	Other:
	The Biodiversity Consultancy; UNEP World Conservation Monitoring Centre; various
	Cambridge academics; Imperial College London
Process and current	The Natural Capital Impact Group is a global network of companies, working collaboratively,
position	to determine how business can sustain the natural world and its resources through its strategies and operating practices.
	Strategies and operating produces.
	Companies in the group are already making progress trying to reduce their impact on
	biodiversity. Companies are supporting the adoption of good farming and conservation
	practices to protect and restore ecosystems. For many of them, the business case for protecting biodiversity is quite simple. Without biodiversity, there is no business.
	protecting area to only to quite an plan in the additional, along to the additional.
	However, leading companies still struggle to showcase their achievements in a manner that
	is consistent, academically robust, easy to understand and comparable. Companies want the opportunity to demonstrate positive impacts and show they are reversing the trend of
	environmental degradation. The Group seek to increase and improve the integration of
	biodiversity into corporate decision making by identifying what to measure and how.
	The Group has developed a concept for a biodiversity metric, recognizing that the largest impacts on biodiversity tend to occur in their upstream value chains, particularly on farms
	where raw materials are produced.
	Development of the Biodiversity Impact Metric commenced two years ago. The development process has included a range of stakeholders at different points in time. There has been
	extensive consultation with stakeholders in the biodiversity conservation community including
	Birdlife International, UNEP WCMC, The Biodiversity Consultancy, Cambridge University
	academics and the Imperial College London. Whilst all members of the Natural Capital
	Impact Group have supported the development of the metric, the global luxury group Kering have championed this effort and provided significant input to ensure that it will inform
	decision making.
	The Beta version of the methodology was launched in July 2018. The methodology is
	currently being piloted with members of the Natural Capital Impact Group.
	CISL is also developing metrics for soil and water. The long-term goal is to produce a metric
	that can be used to assess and track the impact of a company's land use activities on
	biodiversity, soil and water in a given area. Collectively these three elements make up what
	we refer to as the 'Healthy Ecosystem Metric'.
Key features	Biodiversity is inherently multidimensional encompassing taxonomic, functional phylogenetic,
of methodology	genetic, landscape and many other elements of variability of life on Earth.
methodology	It is not practical for a company to measure impacts on biodiversity at all levels in all places,
	so biodiversity assessment methods must be optimized to the specific level of organization
	and spatial scale of interest. Given the multiple variables, the challenge is to narrow the
	scope and identify which variables or proxies can be used for assessing patterns and processes of biodiversity.
	processes of blodiversity.







FICHE NR 2 Biodiversity Impact Metric

The central idea was to develop a robust methodology that is practical, easy to use, and that will help companies to measure their impacts on biodiversity in different parts of the world and provide them with an indication of where these impacts can be reduced. This new measure has the potential to convey impacts on biodiversity in a manner that hasn't been achieved previously and will provide actionable information on how a business can reduce their impact.

The methodology has been co-developed by a number of different companies representing different industries. There is no shortage of targets and indicators that measure biodiversity. However, many of these are difficult for companies to aggregate over all their operations and value chains.

The methodology was developed with the view that whilst biodiversity is complex, business need to report on their biodiversity impacts in a simple way if progress is going to be made incorporating biodiversity into decision making. The methodology for the metric was co designed in line with the following principles:

Metric Principles:

Principle	Description
Meaningful	Meaningful to business and investor communities so it can be used to drive decision making. Methodology is clearly understood.
Measurable and comparable	Allows for comparison across geographies and time.
Possible to aggregate	Can be aggregated from site-level to regional and global scales.
Practical	Data is accessible, measurable by company or using free, globally available data. Ability to substitute better information where available.
Replicable and credible	Based on a reputable scientific method.
Context based	Considers local conditions/levels to reflect 'impact' (beyond 'usage').
Responsive	Responds to changes in company activities, both short and long term.

Earlier working papers providing useful background information are:

- Working Paper on Biodiversity and ES in EP&L
 (https://www.cisl.cam.ac.uk/publications/publication-pdfs/BESinEPLWorkingPaper.pdf)
 (Oct 2016). This work was undertaken to support Kering in developing its thinking on biodiversity and ES in the context of Environmental Profit & Loss (EP&L) accounts.
- Working Paper 'How businesses measure their impacts on nature'
 (https://www.cisl.cam.ac.uk/publications/publication-pdfs/how-businesses-measure-their-impact-on-nature.pdf) (2016)
- Working Paper 'Healthy ecosystem metric framework: biodiversity impact' (https://www.cisl.cam.ac.uk/publications/publication-pdfs/healthy-ecosystem-metric-framework.pdf) (May, 2017)

Metric

Whilst there is never likely to be agreement on an approach that can represent the full spectrum of impacts on biodiversity, there is an urgency to develop indicators that, at a minimum, allow business to see if their decisions are conserving biodiversity or leading to its degradation and loss. Without this knowledge it is not possible to know if the actions being







FICHE NR 2 Biodiversity Impact Metric

taken are working and should continue or if different approaches need to be tried. The biodiversity metric demonstrates that it is possible to show which way some of the key components of biodiversity are heading and what a company can do to change this.

Complex, time-consuming field based biodiversity impact assessments fail to deliver information quickly enough to aid decision-making, particularly for multinational businesses that source hundreds of different raw materials from across the planet.

The Biodiversity Impact Metric is *quantifiable* measure that can be used to *assess* and *track* the impact of a company's *land use* activities on biodiversity in a *given area*.

In essence the metric has been designed to provide information on the following:

- What is the state of biodiversity in an area? How has it changed relative to a baseline?
- How *important* is the biodiversity in an area relative to other areas?
- And how much of the change in biodiversity (both negative and positive) can we
 attribute to the land use of a particular company?

The biodiversity impact measured by the metric relates to the impact of using land that was once natural habitat for commodity production.

The metric combines' data on the land area required to supply a particular raw material with a series of coefficients that quantify the impact on biodiversity.

The basic framework for the metric is as follows:

Biodiversity impact = land area x quantity impacted x quality impacted

Area (ha) of land needed to produce commodity

Company data on amount of commodity and source location

Data on country level yield

Proportion of biodiversity lost through production

Mean species abundance values for land-use types

Relative global importance of the biodiversity lost

Global datasets on range rarity and commodity production by country ecoregion component

The metric aims to characterize the impact of businesses on biodiversity by weighting a company's land area according to its effect on the proportion of biodiversity lost through production (quantity) and the relative importance of the biodiversity lost (quality.) The methodology provides a basis for comparing different sourcing options and as such may help a company to compare different investment options.

Land use impact coefficients are derived using the best available global data to provide a measure of the status of biodiversity stocks and the relative importance of those stocks in a given region. These components are assessed at the eco region level and aggregated to produce a score for different raw materials. The Biodiversity Impact Metric has focused initially on the biodiversity impacts of land-use in relation to raw material supply chains but may have wider application in future.







FICHE NR 2	Biodiversity Impact Metric
Drivers of biodiversity loss	The Natural Capital Impact Group has developed a concept for a biodiversity metric, recognizing that the largest impacts on biodiversity tend to occur in their upstream value chains, particularly on farms where raw materials are produced. The methodology is focused on measuring the impacts of raw material production, or land use, in global supply chains. The biodiversity metric can in principle be used to provide a measure of impact beyond agricultural land use and include extractives and land use in the built environment. The metric will gradually be expanded over time to provide a measure of all impacts on natural capital across a supply chain (i.e. manufacturing/processing/distribution, retail etc.)
Application	
types	In its current form, the metric has particular value for companies involved in bringing a product or service from raw material to market in global supply chains. It supports decision making at a corporate level, providing a commodity-level assessment of biodiversity impacts and giving an indication of where and how a company can reduce their impact in specific places. Raw materials could include raw latex, cotton, rice, coffee, cattle, soybeans, palm oil and other major agricultural commodities.
	This includes companies from the following commercial sectors:
	 Food, beverage and apparel Cosmetics and pharmaceuticals Forest based industries Other major consumer goods companies sourcing raw materials
	The Biodiversity Impact Metric has a number of potential uses, including:
	 Establishing an overall potential impact score from commodity sourcing Flagging geographic sources of potential elevated impacts in a commodity supply chain
	 Comparing potential impacts of different commodities – within a company's supply chains or more generally
	 Comparing potential impacts of different companies sourcing the same commodity(ies)
	The metric is designed to provide robust information on the impact of raw materials that are sourced from different locations across the world. It supports decision making at a corporate level, providing a commodity-level assessment of biodiversity impacts and giving an indication of where and how a company can reduce their impact in specific places.
	Currently the methodology is focused on measuring the largest impacts on biodiversity that occur upstream in most company's value chains. The methodology provides a basis for comparing different sourcing options and as such may help a company to compare different investment options.
	Anglian Water, a water utility company based in the UK, have successfully adapted the methodology to provide an understanding of the current state of biodiversity across all of their operational sites and to prioritize areas where the need to enhance protective measures is more crucial. The methodology will gradually be expanded over time to include different forms of land use and widening the range of companies for whom the metric will be useful.
Required data	Company land area







FICHE NR 2 Biodiversity Impact Metric

Land area is the first data input in the metric. A company provides data on how much land they, or their suppliers, are using to produce a raw material in a particular sourcing location. There are likely to be many cases where a company has limited knowledge of how much land is being used in specific region to produce a specific crop. If this proves to be the case, estimates of land area can be generated using freely available data.

The proportion of biodiversity lost

Biodiversity is inherently multidimensional encompassing taxonomic, functional phylogenetic, genetic, landscape and many other elements of variability of life on the Earth. Measuring change in biodiversity is multi-faceted and can include loss of quantity (abundance, distribution), quality (ecosystem degradation) or variability (diversity of species or genes) within all levels and aspects. It is not practical for a company to measure impacts on biodiversity at all levels in all places, so biodiversity assessment methods must be optimized to the specific level of organization and spatial scale of interest.

Given the multiple variables, the challenge is to narrow this down and identify which variables can be used for assessing patterns and processes of biodiversity. Species diversity is one of the most widely adopted metrics for assessing patterns and processes of biodiversity. Species diversity is strongly correlated with diversity at other levels or organization, such as genetic diversity and ecosystem functioning. The number of different species that are present in a given area is an important measure of ecosystem health and this is often dependent on what type of land use is being employed (e.g. forestry, cropping, pastoral) and how intensively the land is being managed (e.g. monoculture, agro forestry, agro-pastoral)

A series of global coefficients have been derived for different land use types and intensities, representing the amount of biodiversity (in undisturbed natural habitat) that is lost through transformation to a different land-use and land-use intensity. These coefficients that quantify the loss of biodiversity are based on leading empirical studies and provide a basis for differentiating the impacts of land-use types and associated land-use intensities. The values provide an indication of changes in Mean Species Abundance (MSA). MSA is defined as "the mean abundance of original species in disturbed conditions relative to their abundance in undisturbed habitat, as an indicator of the degree to which an ecosystem is intact" (Schipper et al. 2016). These values are based mainly on the latest MSA coefficients (Schipper et al. 2016) with some limited interpolation and expert judgement. Types of land use and management practices carried out by a company across different sourcing locations are categorized according to six broad land cover classifications under minimal, light or intense management.

The importance of a location for biodiversity

All biodiversity is important however in some parts of the world biodiversity is more vulnerable or at risk than in others. Some species are unique to a defined geographic location or have seen their habitat shrink to a point where they are now a great risk of becoming extinct. These species often find themselves marooned in only one, small geographic area and as such are priorities for global conservation. To determine 'biodiversity importance' a new range rarity approach has been developed and incorporated into the metric, drawing on IUCN RED list data. Range data from the IUCN Red List (2017) was used to create a range rarity layer to be used as a proxy of biodiversity importance.

Range rarity values provide an indication of the relative importance of a place for biodiversity compared to other areas. Range rarity scores combine measures of range-restriction (endemism) and richness, considered to be key factors that relate to biodiversity importance. Range rarity is crucial for considering species with very small ranges that are often of greatest conservation concern and is a close proxy for the irreplaceability of a location when the goal is to conserve as many species as possible. Range rarity is determined using Extent of Occurrence range maps for four taxonomic groups that are completely assessed on the








FICHE NR 2 Biodiversity Impact Metric

IUCN Red List of Threatened Species, i.e. amphibians, mammals, birds and conifers. Range rarity has been calculated across ~1 km grid cells. Each ~1km grid cell was scored for range rarity for each species (the proportion of the species' global range the cell represents; i.e. area of grid cell/range size) and given a total score by summing scores across all the species potentially occurring in it.



Visualization of range rarity by Country Ecoregion Component (analysis and mapping: UNEP-WCMC)

Areas that hold a large number of species (richness) and/or species with small ranges sizes are given a higher score. The species in these areas are more vulnerable to land conversion. The taxonomic groups used to represent biodiversity are limited to amphibians, birds, mammals and conifers. Other taxonomic groups were not considered for this analysis as they are still incomplete, at least for the terrestrial realm, and the available spatial data will typically have been collated with a number of geographic biases. As the IUCN Red List assessment and mapping process is ongoing, further groups could be included in future leading to provide a more representative index of biodiversity importance. The global pattern of importance based on aggregation to the country-ecoregion combination (CEC) level shows expected patterns of high importance in the tropics and especially on small islands and in mountain ranges.

Calculating an impact score

The metric can be applied in different ways to calculate an impact score, depending on the amount of information a company is able to provide on its sourcing activities:

- 1. Tier 1: Calculating a 'base' score (with limited company sourcing information)
- 2. Tier 2: Calculating an 'improved' impact score (with information on a commodities source location, production management and local yields)

Tier 1: Calculating a 'base' score (with limited company sourcing information)







FICHE NR 2 **Biodiversity Impact Metric** At this level the metric can be used to identify potential hotspots or risks in a supply chain associated with the production of particular commodities. It can provide an indication of the magnitude of the impact they are having and where there are opportunities to reduce their impact through their operational decision making. The metric can be used to: Flag geographic sources of potential elevated impacts in a commodity supply chain Compare potential impacts of different commodities – within a company's supply chains or more generally Comparing potential impacts of different companies sourcing the same commodity(ies) The metric can be applied at this level using information on the amount of production sourced from a particular country, or region. To calculate the metric in its simplest form, the input data required is the weight of the commodity sourced (in tonnes) from each producing country. The metric calculates the biodiversity impact of a company's commodity sourcing in weighted hectares (wehas.) A weighted hectare combines land area (hectares) with weightings for biodiversity importance, based on species range rarity, and quantity of biodiversity impacted, based on global averages for Mean Species Abundance. In essence, what this means is that each hectare of land under production is given a weighting, providing an indication of how much biodiversity has been lost as a result of land conversion and how important the loss of that biodiversity is relative to other areas. This impact can be expressed as a total (i.e. 320 weighted hectares) or as an averaged impact/tonne. Impact scores can be compared across different locations or against a global average to provide an indication of whether the sourcing of a particular commodity is having a below or above average impact on biodiversity in a particular area, compared to other sourcing locations. At this level, the metric distinguishes between quantity of biodiversity impacted by different land-use types, but not by different land-use intensities. Tier 2: Calculating a more refined impact score accounting for different land-use intensities in a specific geographic location The relationship between biodiversity and how intensely land is managed is complex, however low intensity production systems tend to harbor more biodiversity than intensive systems. For focusing and tracking improvements in land management and production practices, the 'detailed' version of the metric is required. At this level the metric can be used to start distinguishing between the impacts of different land-use intensities. If more detail is available on source location, production management, and local yields, then the metric may be applied at a 'detailed' level, with land-use intensity also be factored into the metric. At this level a company can start to identify the links between different supply chain interventions (internal standards, third party certification schemes) and reducing impacts on biodiversity in different location. **Granularity** The various components in the metric are assessed at the eco region level. Terrestrial level Ecoregions of the World are a biogeographic regionalization of the Earth's terrestrial biodiversity. Ecoregions are defined as relatively large units of land containing a distinct assemblage of natural communities sharing a large majority of species, dynamics and environmental conditions. The new terrestrial ecoregion dataset²⁵, updated in 2017,

proposes 867 terrestrial ecoregions, which size ranges from 3 km² (Central Polynesian






28






FICHE NR 2 Biodiversity Impact Metric

tropical moist forests) to 3.9 million km² (East Siberian taiga), with an average size of ecoregions of 155,748 km². They are classified into 14 different biomes such as forests, grasslands or deserts.



Ecoregions are a convenient size, small enough that ecosystem type and biodiversity 'quality' are largely consistent within them, but large enough that precise source information for commodities is not needed. Most ecoregions tend to stretch across national boundaries.

A 'Country Ecoregion Component', or CEC, (see map below) is the portion of a particular ecoregion that falls within a particular country. The map below provides an example of different Country Ecoregion Components in Liberia.



Metric results are calculated at the scale of Country Ecoregion Component and then aggregated to produce a national score for different commodities. The metric can work at any scale in principle, but the underlying information base has been developed at the scale of Country Ecoregion Component.

Strengths recognised by initiators

- The metric is academically robust and draws on a number of globally recognized data sets. It does not involve primary, field based data collection in different sites
- It is simple to use and doesn't require a company to provide large amounts of data
- It provides a comparable methodology to showcase progress across different industries
- There are a number of steps or stages in Natural Capital protocol, starting with a framing and scoping phase and then progressing into measurement, valuation and







FICHE NR 2	Biodiversity Impact Metric
	application. The Metric can help companies during the Scoping and Measuring stages of applying the protocol
Weaknesses recognised by initiators	 It will help shed light on high-risk locations where the company is most likely to experience biodiversity risks but won't provide enough information about a specific site to support completion of a biodiversity action plan. This would require more detailed information at a finer scale. At present, it is focused on the impacts of land use associated with raw material production. It does not provide a measure of all impacts on natural capital across a supply chain (i.e. manufacturing/processing/distribution, retail etc.) The metric does not assess the broader landscape context, nor the indirect effects (outside the land-use footprint) of commodity production. The metric does not assess conservation status of commodity-producing ecoregions. Supplementary metrics can be calculated (from ecoregion-scale data) to assess the conservation context of commodity sourcing from a particular country, in relation to the area of ecoregion protected and the extent of unprotected natural habitat remaining.







3. Biodiversity Indicator for Extractive Companies (UNEP-WCMC)

FICHE NR 3	Biodiversity Indicator for Extractive Companies
Date of	First assessment by UNEP-WCMC on 2 nd April 2018
assessment	Updates on 15 th July 2018 and 21 August 2018
Actors	Lead: UNEP-WCMC Other: IPIECA Biodiversity and ES working group (http://www.ipieca.org/our-work/environment/bes-issue-management/) and Proteus Partners (https://www.unep-wcmc.org/featured-projects/proteus-partnership) engaged as a platform to provide industry specific insights for the development of an indicator. An advisory group comprising: ConocoPhillips, Conservation International, ENI, IUCN,
	Oxford University, Rijksdienst voor Ondernemend Nederland (RVO.nl), Shell, Fauna & Flora International and The Biodiversity Consultancy provide an independent sounding board to the project.
Process and current position	UNEP-WCMC initiated the Biodiversity Indicator for Extractive Companies project in March 2017. Phase 1 determined the needs and current practices used by the sector, through desk review and interviews with industry representatives. The key findings of Phase 1 highlighted two potential models to take forward for further methodological development and piloting: A single composite indicator A framework approach
	For a detailed description of the findings of phase 1, please find the report here .
	Phase 2 of the project commenced in January of 2018 (currently ongoing). A shortlist of suitable methodologies, that fit the models described above, was created and explored with industry representatives at a half day workshop. Participants highlighted that a single metric may not provide the necessary resolution, flexibility or information to meet the indicator needs of the extractive sector. Current data gaps and lack of comparability may in the short-term render a single indicator challenging.
	Learning from Greenhouse Gas reporting, a tiered approach, incorporating high-level screening of all sites within a portfolio followed by the development of indicators for identified high-risk sites, using a Pressure-State-Response model were identified as a potential first step in the development of improved biodiversity indicators for the extractive sector and would draw from existing methodologies and data.
	A draft methodology was discussed at the Proteus Annual Member's. It was positively received by IPIECA and Proteus members. The methodology is now being finalized following this feedback and Phase 3 of the project is being developed to pilot the methodology.
Key features of	A three stages process is suggested for indicator development (
methodology	Figure 1):
	 First stage: screening of the company's portfolio of operations to identify sites with potentially high biodiversity sensitivity, based on globally and locally available data sets combined with site validation to identify sensitive sites for prioritisation of reporting effort;
	Second stage: development of site level biodiversity indicators using the state-pressure-response (SPR) framework (a widely accepted organising framework for biodiversity management and monitoring), informed by the stage above and based on site level data and documentation for high sensitivity sites collected as part of the environmental impact assessment; and







 Third stage: aggregation of scores for SPR at site level up to business unit, division, and corporate level to provide insight into performance on the ground.

Figure 1: Process for site prioritization and biodiversity indicator development

Second Stage First Stage Third Stage Applying site level **Biodiversity** Aggregation and state pressure response framework sensitivity screening reporting Identify site level metrics against framework Screen to identify high sensitivity sites Validate results with Calculate Reporting and disclosure Aggregate site SPR site dashboard **Expected Output Expected Output Expected Output**

 A final list of all sites categorised by potential biodiversity sensitivity (low / medium / high)

- A state pressure response score for each site
- A site indicator dashboard
- Aggregated corporate SPR scores
 - Corporate-level indicators / disclosure

Metrics

In order to comply with the user needs of consulted companies, the following criteria apply to the envisaged indicator(s).

Criteria	Description
1.Business relevant	Indicator can be used directly for company's business decisions at multiple levels.
2.Spatial extent	The ability to use an indicator at site level and aggregate it up to business unit or corporate level.
3.Sensitivity to change	Indicator responds to change in the issue of interest with minimal lag, enabling monitoring over time.
4.Data availability /accessibility	Extent to which data is available and inexpensive/ feasible to access.
5.Scientifically valid*	Accepted theory of relationship between the indicator and its purpose, with agreement that change in the indicator indicates change in the issue of concern.
6.Communication	Ease of understanding of indicator to non-technical people.
7.Production	Data collection, analysis and calculation is straightforward and non- specialists can use the indicator.
8. Policy relevance*	Links to SDG/Aichi Targets.

The below table provides an overview of screening criteria, and recommended data source for global-scale screening (First Stage in Figure 1 above).







	Criterion	Description	Data source
	Criterion 1: Globally threatened species	Criterion evaluating the number of threatened species ranges overlapping an operating site, taking into account the size of the ranges. Operating sites are scored based on overlap with a range rarity layer for threatened species, derived from the IUCN Red List species ranges. Threatened species are species listed on the Red List as Critically Endangered (CR), Endangered (EN) and Vulnerable (VU). Range rarity measures species richness weighted by the inverse of range size, giving high values to locations that support more species that are found in few other locations. For the purposes of this criterion, range rarity is further weighted by relative extinction risk, giving higher scores for locations support species that are more threatened.	Red List spe range rarity I 1x1km resol (IUCN). Avai through IBAT of January 2
	Criterion 2: Critical habitat	Criterion evaluating the overlap of operating sites with areas which likely or potentially classify as Critical Habitat, as defined by the IFC Performance Standard 6 (PS6) (IFC 2012). IFC PS6 defines critical habitat as areas of high biodiversity value, based on a set of five criteria: • Habitats of significant important to Critically Endangered and/or Endangered species • Habitat of significant important to endemic and/or restricted-range species • Habitat supporting globally significant concentrations of migratory species and/or congregatory species • Highly threatened and/or unique ecosystems • Areas associated with key evolutionary processes As detailed in the IFC performance standard guidance note 6, Critical Habitat may also be triggered by other recognized areas of high biodiversity values, including Protected Areas (Ia-II and sometimes III-IV), World Heritage Sites and the majority of Key Biodiversity Areas, which encompass Important Bird and Biodiversity Areas and Alliance for Zero Extinction sites.	Global Critic Habitat scre layer (UNEP WCMC). Ava through IBAT of January 2
	Criterion 3: Protected areas	Criterion evaluating the overlap of operating sites with national-level protected areas and protected areas designated under regional or international conventions or agreements. Protected areas aim to conserve biodiversity by protecting species, habitats and other biodiversity features within their boundaries. Protected areas are one of the cornerstones of biodiversity conservation as outlined by the Convention on Biological Diversity (CBD). The authoritative definition of protected areas has been formulated by the IUCN (Dudley 2008).	Polygon-bas World Datab on Protected Areas (IUCN UNEP-WCM Available thr IBAT.
Outcomes based on modelling or real data?	It is intended that the approach uses real data, but modelled/ estimated data may also be used. This will be explored within the piloting process.		_
Drivers of biodiversity loss	Pressures are considered as a part of the application of the state, pressure, response model.		
Application types	Indicator for applications:	ronment World Conservation Monitoring Centre (UNEP-WCMC) Bir Extractive Companies can support the following types of business essing and monitoring progress of biodiversity risks at project, site a corate level aparing options at project and site level ying a NNL approach at project, site and corporate level (establishing monitoring progress)	nd







	 Communication and reporting (e.g. compliance with GRI reporting) Both new and existing operations FIs can compare biodiversity performance of companies within the extractives sector 		
Required data	For Stage 1: Data on critical habitat, IUCN red list species range rarity information and location of protected areas, derived from the Integrated biodiversity assessment tool (IBAT). The initial screen looks at globally threatened species as a quick and easy set of data to access, this creates a short list of potentially sensitive sites. For Stage 2: Those priorities are then checked through site discussion and with time all sites will be reviewed. IBAT is not the source of information for the site level indicator production, EIAs, BAPs (biodiversity action plans) and local data sets will be.		
Granularity level	It will cover scales at both the site level and corporate level.		
User friendliness	To be determined through piloting. Initial feedback on the concept has been positive.		
Strengths recognised by initiators	 Builds from existing environmental management systems Stage 1 formalises and documents existing industry practice enabling a more transparent communication of identification and management of risks Designed in consultation with industry and driven by industry needs Builds from a model well used in the conservation and policy arena Theoretically sensitive to management interventions 		
Weaknesses recognised by initiators	 Assessment looks at current status of biodiversity rather than future predictors of decline Incompleteness of data sets may lead to understatement or overstatement of sensitivity Isolating pressures attributable to corporate action is challenging, as is defining area of influence The resultant indicator set may be complex to communicate and may not meet needs for external disclosure Lags in responsiveness of indicators may obscure performance 		







4. Product Biodiversity Footprint (I Care & Consult, Sayari)

FICHE NR 4	Product Biodiversity Footprint		
Date of	First version by I Care & Consult on 20 June 2018		
assessment	First review EU B@B Platform on 6 August 2018		
	Update by I Care & Consult on 23 August 2018		
Actors	<u>Lead</u> : I Care & Consult, Sayari		
	Other:		
	Steering committee: French ministry of environment, ADEME, Kering, Avril, L'Oréal		
	Scientific committee: members of the following organizations: MNHN, UN Environment, IUCN, Irstea, Solinnen.		
Process and	The PBF project aims to answer the lack of specific tools to assess the impact of		
current	different products and services on biodiversity. In order to do this, baseline principle of		
position	the PBF project is to co-develop a method and a tool relying both on sectoral/local		
	biodiversity studies and on company data to quantify the impacts of a product on		
	biodiversity all along the product's life cycle stages in order to provide		
	recommendations for changes.		
	PBF project brings together all existing available data and provides quantitative results		
	for decision making processes regarding product strategy (risks analysis, purchasing		
	strategy, eco-design). The approach for development of the first version is presented in a method document,		
	available on the project website: http://www.productbiodiversityfootprint.com/ . It has		
	intrinsically combined scientific knowledge (embodied by the scientific committee,		
	gathering experts from both LCA and ecological field) and business requirements and		
	case studies.		
	Product Biodiversity Footprint is about bridging a gap between 2 worlds : field		
	analysis on one side, lifecycle analysis based on data on the other side		
	Life Cycle Initiative		
	RED ISSG protected planet net		
	LIST Specialist Group		
	CountrySTAT Rod and agriculture data activotic		
	Food and agriculture data network Food and agriculture data network Food and agriculture data network Food and agriculture data network Food and agriculture data network Food and agriculture data network Food and agriculture data network		
	f tomiquation formation		
	GIBIO Introductional processing and the processing		
	Agocidating human impacts on biodiversity Agocidating human impacts on biodiversity Tail impairs (Date Clings Tail impairs (Date Clings) (Date Clings)		
	16 there comp 16 there is a compared to the co		
	The ecologists world The LCA practitioners world		
	First assessments took place in 2017, and 3 case studies provided results in February		
	2018 for three agricultural sectors: goat wool, rapeseed oil, palm oil (phase 1). Further		
	testing for other sectors will take place in 2018- 2019 (phase 2)		
Key features of	Discriminating capacity		
methodology	The main objective of the project is to improve the biodiversity performance of a product		
	by identifying biodiversity hotspots that can be improved and support eco-design approaches. In order to have such a capacity the PBF has a strong discriminating		
	capacity: the method aims to identify between the variants of a product the one with		
	lowest impacts on biodiversity.		
	The LCA framework is used to calculate the relative differences between the variants		
	of a product. In the long term, it will also allow to compare different products or different		
	sectors at larger scales.		
	Internating his diversity in LCA acceptation		
	Integrating biodiversity in LCA ecosystem		







FICHE NR 4 Product Biodiversity Footprint

To be easily adopted by companies for product assessment, the method is integrated in the LCA ecosystem, meaning it is **connected to LCA database** and compatible with **LCA assessment framework** (Chaudhary et al. 2015, Verones et al. 2017)²⁶., so that this additional biodiversity assessment can be seen as an **add-on to LCA global assessment**. Therefore, the choice made for PBF method is to include biodiversity knowledge in the LCA framework.

Covering a large scope of impacts

The method covers the 5 pressures on biodiversity identified in the Millennium Ecosystem Assessment (2005). Biodiversity knowledge included in the LCA framework is based on **ecological publications** specific for each pressure and on **available global biodiversity database** to assess the state of biodiversity.

Taking into consideration vulnerability and irreplaceability of species

Metrics

PBF allows the comparison of different product scenarios: one "reference" and one or more "variants".

The comparison is made on an indicator reporting the **potential loss of species**. The indicator is expressed in the so called "potential disappeared fraction of species within a year" or PDF*yr. This indicator is developed by LCA researchers, and frequently used in LCA methodologies. It can be applied both as a regional and global indicator, as recommended by the UNEP SETAC 2016 guidance:

- Regional indicator (PDF_{reg}*yr) quantifies the potential for disappearance of species at regional level; region is understood as an ecologically homogeneous area, practically identifies as ecoregions for terrestrial ecosystems, and the water basins for freshwater ecosystems).
- Global indicator (PDF_{glo}*yr) quantifies the potential for global extinction of species, accounting for their vulnerability at global level.

Results are presented in different layers (see figure below):

- (a) A first level displays a relative comparison of the biodiversity impacts on the 5 axes of MEA pressures,; the 100% value will be assigned to the Reference scenario;
- (b) The second level provides details for each MEA pressure: relative comparison and absolute quantification.
 - Change of habitat is detailed in
 - land occupation impact on biodiversity
 - land transformation impact on biodiversity
 - water stress impact on biodiversity
 - o Pollution is detailed in:
 - terrestrial acidification impact on biodiversity,
 - freshwater eutrophication impact on biodiversity
 - photochemical oxidation impact on biodiversity
 - Climate change impact on biodiversity is displayed in biodiversity loss. Impacts will also be supplied in the usual unit of kgCO2eq, according to IPCC 2013 100-year factors (In Module 1, the relationship between pressure and impact on biodiversity is modelized following LC methodology (2016), as precised in the footnote. Module 2 helps refining these impacts through local data and details on company practices).


²⁶ Verones et al. 2017. LCIA framework and cross-cutting issues guidance within the UNEP-SETAC Life Cycle Initiative, *Journal of Cleaner Production*, vol. 161. Chaudhary et al. 2015. Quantifying Land Use Impacts on Biodiversity: Combining Species– Area Models and Vulnerability Indicators, *Environ. Sci. Technol.* 2015, 49, 9987–9995.



_








PBF two main outputs: two different levels of results aggregation, to assess the biodiversity impact of two variants of one product (fictive product).

To calculate these metrics, the methodology is based on 3 modules, as described in figure bellow:

Module 1 computes lifecycle impact assessment, with spatial differentiation for the main impact categories evaluated. Characterization factors used for computation are the ones currently available in published LCA methodologies (LC Impact²⁷). This first module enables the user to visualize the hotspots of the product footprint both geographically and along the whole value chain.

- Module 2 treats specific information regarding the practices and the local context, that would enable to adjust impact computations based on information entered by the users. The development of criteria and rules to quantify the changes in impact computations are defined per type of land use (e.g. arable crops, perennial crops, grassland, mining, forest, urban....) and/or per sector (livestock, transport, electricity, construction....). Results of modules 2, additionally to the results of the Module 1, enable the user to visualize and quantify the benefits of a chosen practice/location, and compare various scenarios for a given product.
- Module 3 assesses qualitatively 2 aspects that are not part of any LCA model, namely 'invasive species' and 'species management', this last one encompassing 'overexploitation' aspects of the Millennium Ecosystem Assessment, such as hunting, poaching or overfishing, but going beyond as it also includes positive actions (e.g. installation of pollinators, use of various breeds, follow up of endangered species...).









FIGUE ND 4	Breakert Biodiversity Footpuint							
FICHE NR 4 Outcomes	Product Biodiversity Footprint Outcomes are based on a mixed approach, with modelling and real data.							
based on	Outcomes are based on a mixed approach, with modelling and real data.							
modelling or								
real data?								
Drivers of	Biodiversity impacts are assessed independently for the 5 MEA pressures: habitat							
biodiversity	change, climate change, pollution, species management and invasive species. At							
loss	this stage, we did not combine the 5 pressure scores into a single biodiversity score, to							
	avoid issues of scales and weighting between indicators.							
Application	 Product evaluation: PBF evaluates the impact of products or services created as a result of a fabrication, manufacturing, or production process. This approach is mainly based on life-cycle assessment (LCA), a technique to assess environmental impacts associated with all the stages of a product's life. PBF can help avoid a narrow outlook on environmental concerns and be a tool for decision-making or product communication (see below). Decision making Sourcing of material Change of production practices (e.g. farming) Eco-design 							
	 Product communication: the result of the comparison between reference and variant can help the company in producing convincing and scientific-based points for enhancing and valorising the biodiversity benefits of the product 							
Required data	 Company data: the company needs to transfer following data to PBF team Standard Data on the production process of the product: raw materials, transport, yield, Localization data for the main processes: localization of sourcing, Existing local biodiversity studies (inventories, impact studies,) List of actions undertaken/ practices put in place to mitigate the impact on biodiversity Method data LCI Data (Ecoinvent (https://simapro.com/databases/ecoinvent/), Agribalyse (https://simapro.com/products/agribalyse-agricultural-database/),) Biodiversity database (IUCN, IBAT, Predicts,) 							
Granularity level	 General level: ecoregion and countries. although spatial resolution is specific to each impact category, LC-Impact proposes aggregated characterization factors at country level for each environmental category. Detailed level: land use occupation practices and biodiversity local context (depending on the level of data available on the product) 							
User friendliness	 Required expertise: There is a need for both a LCA expert and a biodiversity expert, to gather the necessary input data for PBF tool, Time effort Time effort is limited, quite similar to a normal LCA (provided that underlying biodiversity data are already existing): 10% ETP during 4 months for LCA specialist and 10% ETP during 4 months for biodiversity specialist Specificity of the approach The approach is actually a mix of other environmental approaches already existing in the company: LCA approach and impact study approach Communication of results Thanks to the spider graphic, it is very easy for a non-expert to understand the results of PBF 							

²⁷ Verones F et al., 2016, LC-Impact Version 0.5, A spatially differentiated life cycle impact assessment approach, retrieved from http://www.lc-impact_eu/downloads/documents/LC-Impact_report_SEPT2016_20160927.pdf







FICHE NR 4	Product Biodiversity Footprint
Strengths recognised by initiators	 The testing of the PBF methodology has revealed 4 main strengths A full "product" approach, encompassing all the lifecycle of the product and not only one specific phase (e.g. agriculture production) Ability to reveal the impact of the positive biodiversity actions of a company along the product lifecycle (sourcing, production practices,) Capacity to combine both "database modelized information" and real company data, so that it reduces the need for input data from the company but also positions the specific product performance vs. average product performance The coverage of all pressures on biodiversity (and not only impact of land use)
Weaknesses recognised by initiators	This methodology is a significant step towards pragmatically assessing the biodiversity impact at product level. Application to case studies has proven the feasibility of the method. Still some weaknesses have been identified and some improvements will be performed in 2018/2019 • Methodology needs to be tested and adapted to other sectors: Energy production, Mining, Transport Infrastructures in particular. • Methodology needs to include or complete some cause-effect pathways, by example adding ecotoxicity in pollution; vulnerability in further LCA categories, • Graphic interface (maps) needs to be improved to facilitate vision of impacts geographically







5. Biodiversity Footprint approach (ASN Bank)


FICHE NR 5	Biodiversity Footprinting approach						
Date of assessment	21 June 2018						
Actors	<u>Lead</u> : ASN Bank (The Netherlands) <u>Other</u> : The methodology is co-developed with CREM and PRé Consultants.						
Process and current position	ASN Bank launched the methodology in 2016. The first footprinting exercise and associated report were made and published in August 2016. In 2017 a second report was published in which a footprint of the whole balance sheet was made over the years 2014, 2015 & 2016. The methodology was also adjusted with this second footprint, based on external and internal methodological and data updates. The method entails both a quantitative and qualitative assessment. At this moment ASN is working together with CDC Biodiversité, ACTIAM and Finance in Motion to establish a common ground for biodiversity footprinting by financial institutions. They will publish a joint paper on this in November 2018.						
Key features of methodology	ASN's Biodiversity Footprinting approach is designed to provide an overall biodiversity footprint of the economic activities a financial institution (FI) invests in. The <u>quantitative</u> methodology consists of 3 steps:						
	 The <u>first step</u> is to create an overview of the economic activities the FI invests in. This step includes: A 'definition' of the activities of a company: what is the company producing (in what sectors is the company active?) and where does production take place? Decisions on the scope of the assessment, like the inclusion of supply chains of companies the FI invests in (included in the ASN Bank approach); A selection of the investments included in the assessment (all major investments) In the <u>second step</u> the environmental impact of the economic activities of the companies invested in is assessed. The environmental data in the 'Exiobase' input-output database²⁸ is used to assess what land use, water use, emissions, etc. is linked to the economic activities of the companies. Exiobase takes into account world-wide trade flows between countries and between sectors. In the <u>third step</u>, the ReCiPe methodology is used to calculate the environmental footprint on a midpoint level (e.g. climate change resulting from CO2 emissions) and to calculate the resulting impact on ecosystem quality or biodiversity (endpoint level). This latter step is based on science based 'dose-response' relations (e.g. the effect of a 1 degree temperature rise on biodiversity). This results in an impact on terrestrial biodiversity and an impact on aquatic biodiversity. The unit used to express the impact on biodiversity is PDF.ha.yr, the Potentially Disappeared Fraction of species per hectare (per cubic meter for aquatic biodiversity) per year. The result is then used to calculate the biodiversity footprint in m2 per Euro invested (for each investment category) and the total footprint in m2 for all investments. A <u>qualitative</u> analysis is used to guide the interpretation and the use of the footprint results, looking at (among others) the limitations of the footprinting methodology and the potential influence of the footprint results on investment decisions. 						

















FICHE NR 5	Biodiversity Footprinting approach						
	 Exiobase data; the use of other data-sources (like Trucost data) is being explored. Information on biodiversity impact drivers in different sectors to allow for a qualitative analysis guiding interpretation and use of the footprint results. 						
Granularity level	Method is best suited to get a broad overview and footprint and to determine the hot spots of biodiversity impact in the portfolio /balance sheet.						
User friendliness	The methodology is user friendly, but requires the use of impact calculation software fit to deal with the (high number of) input data and the calculation of impacts based on the ReCiPe methodology. Depending on the level of knowledge within the FI using the methodology, expert input may be needed to stay informed of data and methodology updates and to enable a correct interpretation of the footprint results.						
Strengths recognised by initiators Weaknesses recognised by	 Scientifically well underpinned Use of open source and free database (no black box calculations) The Exiobase input/output model shows trade flows between countries and sectors and therefore allows for a geographical identification of impact hot spots on a country level. Supported by range of stakeholders (including government, knowledge institutes and NGOs) after stakeholder consultations Covers most drivers for biodiversity loss. Location/region specific data can be used when they are/become available. Scalable to be used by other banks The ReCiPe methodology takes into account pollution (besides nutrient load and other impact drivers) The complementary qualitative analysis guides correct interpretation and use Exiobase limitation (use of sector average data). This weakness is expected to improve and change in time when better data will become available. 						
initiators	 Land-use related impacts are biased to temperate regions which means that land-use related impacts will be less accurate for tropical regions. Inclusion of location specific characteristics is limited, limiting the methodology's fitness for use on a project level. On a portfolio level, with the aim of identifying biodiversity impact hot spots, this limitation is fine. Not all drivers of the loss of biodiversity are covered by the ReCipe methodology. For example, the introduction of invasive species and overexploitation are not covered. This limitation is addressed by means of a complementary qualitative analysis, analysing the significance of this limitation and what this means for the interpretation of results. 						







6. The Biodiversity Return on Investment Metric (by IUCN)

FICHE NR 10	Biodiversity Return on Investment Metric (BRIM)							
Date of assessment	• 29 August 2018							
Actors	Lead: IUCN							
Process and	Other: Vulcan Inc; The Biodiversity Consultancy; Birdlife International Technical approach and methodology developed during 2017 through a series of							
current	consultative workshops and a research program. Methodology under road test in two field							
position	situations- Sumatra and Nicaragua. Process underway to submit manuscript of							
	methodology and lessons learned from road-test in peer-reviewed journal by end 2018.							
Key features of	BRIM measures the change in risk of species extinction attributable to							
methodology	 investment Investment can change the scale and impact of the processes that cause species 							
	Investment can change the scale and impact of the processes that cause species to be at risk of extinction- deforestation, over-exploitation							
	Based on the IUCN Red List of Threatened Species – the global standard for							
	documenting species' conservation status							
	 Based on quantitative categories and criteria – not expert opinion Incorporates compilation of data on range, habitats, threats, etc – not just a list 							
	incorporates compilation of data off range, habitats, tiffeats, etc. – not just a list							
Metrics	A fully additive and scalable metric, from pixel to global, or across sites in a							
	 portfolio BRIM allows comparison across investment targets – a change in biodiversity 							
	value in one place is directly comparable to a change somewhere else on the							
	planet							
	Responsive at the pace of investors; changes in management can quickly cause							
	changes in pressures affecting species							
Outcomes	Based on the IUCN Red List of Threatened Species – the global standard for							
based on	documenting species' conservation status							
modelling or real data?	 Based on quantitative categories and criteria – not expert opinion Incorporates compilation of data on range, habitats, threats, etc – not just 							
	a list							
Drivers of biodiversity	Based on the IUCN/Conservation Measures Partnership (CMP) Unified Classification of							
loss	Direct Threats.							
Application	BRIM focuses on identifying those actions which generate the highest Rol in terms of							
	biodiversity value. So, it's <u>not a financial tool</u> . The cost of the measures is not included. It is a separate component.							
	Bank: Screening of potential biodiversity ROI across a portfolio (for instance,							
	investments in a particular company value chain, for a site-based commodity)							
	National Government: National baseline of potential for species extinction							
	reduction, establishment of SDG 14 target and progress measures towards targ							
	 (contribution to change in Red List Index) Conservation project investor: ex-ante evaluation of project potential to reduce 							
	risk of species extinction; identification of priority management actions; ex-post							
	progress towards target							
	Company: assessment of potential for species extinction risk reduction across							
Required data	 corporate footprint (mine sites, plantations, landscape restoration) Species range maps or maps of Extent of Suitable Habitat from the IUCN Red 							
Toquii ou data	List of Species							
	 Threat status and threats applying to species from the IUCN Red List of Species 							
	(complemented by information from the Key Biodiversity Areas dataset)							
	Polygon of investment intervention is needed (site, protected area, country, landscape, land use unit)							
	landscape, land-use unit)Site- or country-based measures require bespoke assessments using GIS							
	Site- or country-based measures require bespoke assessments using GIS applications and integration of threat and land use layers							







FICHE NR 10	Biodiversity Return on Investment Metric (BRIM)					
Granularity level	Measurable at any scale from pixel to global. Additive and scalable, and comparable globally, so that changes in BRIM in two sites in different parts of the world are directly comparable					
User friendliness	 Is it easy to use by non-experts or does it require specialist knowledge? The methodology can be interpreted by non-experts and recommendations made to managers based on results of assessments Site- or country-based assessments require bespoke assessments using GIS applications and integration of threat and land use layers, by technicians familiar with the IUCN Red List of Species What are the required time efforts? Under assessment Does it require a similar approach as other environmental issues or is it very specific? Species are the best-known component of biodiversity Their status has been evaluated in a scientifically consistent, multi-stakeholder, global process (the IUCN Red List of Species) The presence of threatened species in a site or habitat is an indication that the ecosystem is under pressure Habitats or ecosystems with many threatened species are likely to be at a greater risk of collapse Ecosystem collapse will cause significant harm to people and economies Investing in reducing the risk of species extinctions is an efficient way to maintain ecosystem benefits to people 					
Strengths recognised by initiators	 It can help the finance industry and investors target their investments to achieve conservation outcomes BRIM can enable investors and governments to track conservation gains BRIM can measure the contributions these investments make to global targets such as the Sustainable Development Goals, as it can be used to track progress towards changes in the Red List Index Based on the IUCN Red List of Threatened Species – the global standard for documenting species' conservation status Based on quantitative categories and criteria – not expert opinion Incorporates compilation of data on range, habitats, threats, etc – not just a list BRIM allows comparison across investment targets – a change in value in one place is directly comparable to a change somewhere else on the planet A fully additive and scalable metric, from pixel to global, or across sites in a portfolio Responsive at the pace of investors- changes in management can quickly cause changes in pressures affecting species The BRIM can measure ex-ante (potential) and ex-post (achieved) impacts of investments at a range of scales and over a range of timeframes. 					
Weaknesses recognised by initiators	 It can complement conventional measures of financial ROI Not all species are adequately scored for extent and intensity of threat-full calculation of BRIM will require assessment for some or many species Not all taxa are comprehensively assessed (some species have not been evaluated at all, coverage in some ecosystems, such as marine and freshwater, is incomplete) Application at site scale requires some data gathering and complementary information Measurement of ex-post BRIM requires identification of linkages between investment and changes in correlates of population (for instance changes in Extent of Suitable Habitat) 					







7. The Agrobiodiversity Index (ABD) by Bioversity International

FICHE NR 6	THE AGROBIODIVERSITY INDEX (ABD index)						
Date of	First version by Bioversity Int. on August 1, 2018						
assessment	First review by EU B@B Platform on 7 August						
Actors	Lead: Bioversity International Other: Clarmondial AG, a Swiss investment advisory company focused on sustainable natural resources investments is a partner for business and finance applications. Bioversity International is a member of the World Benchmarking Alliance platform, which aims to provide information that indicates how companies are contributing to the SDGs and is helping to leverage and harmonize SDG-related monitoring initiatives under development. Several knowledge and data partners contribute to the development of the ABD index including, among others: FAO, the PREDICTS group of the Natural History Museum and the Commonwealth Scientific and Industrial Research Organization (CSIRO). EU DEVCO						
	and Italian Cooperation are pioneering funders.						
Process and current position	 Process: A feasibility study mapped the demand and specific needs for an Agrobiodiversity Index among food and agriculture companies, institutional investors, governments as well as other index groups. A review of the scientific literature was published and provides the scientific foundations of the Agrobiodiversity Index, in particular the importance of agrobiodiversity for food 						
	 system sustainability, healthy and diverse diets, production systems resilience to abiotic and biotic stresses, and for seed systems, agroecological intensification, ES and conservation. A first prototype was developed and an online portal created (completed June 28, 2018) (https://www.bioversityinternational.org/abd-index/) The prototype has been populated with data from existing global databases and from case studies from pioneering companies and countries. This brings, for the first time, several datasets together from an agrobiodiversity perspective. A sensitivity analysis was performed to assess different ways of aggregating data. An advisory panel to guide the use and scaling of the ABD index has been established, including relevant experts from business alliances, multi-lateral government bodies, civil society and academics. Continuous interactions with private and public sector users feed into the design and applications of the ABD index. Current position: The ABD index prototype Version 1 and the related online portal are ready and will be officially launched in November 2018. A process to continuously strengthen the ABD index, integrate new data and fill specific 						
	 A process to continuously strengthen the ABD index, integrate new data and hir specific data gaps, is being set up. Food and agriculture company and country case studies and applications are being elaborated. The companies represent different actors along the value chain – input suppliers, processors, retailers. The countries represent different continents and agroecological systems. Continuous interactions with private and public sector users feed into the fine-tuning and applications of the ABD index. 						
Key features of methodology	The ABD Index fills a specific niche – on agricultural biodiversity the foundation of agroecological intensification and food system sustainability, which is fundamental for both food and agricultural companies and countries, to reduce material risks, seize new market opportunities, and make global food systems sustainable.						
	The focus is on agricultural biodiversity at the genetic, species and landscape levels, i.e. the variety and variability of animals, plants and micro-organisms that are used directly or indirectly for food and agriculture (FAO definition). This makes the ABD index very complementary to other metrics or indices that focus primarily on wild biodiversity. Research shows many interdependencies and linkages between						








THE AGROBIODIVERSITY INDEX (ABD index) FICHE NR 6 agrobiodiversity and wild biodiversity. Both wild biodiversity and agrobiodiversity provide multiple ES that support food production, underpin food security and human well-being. A review for the UN Convention on Biological Diversity of 119 National Biodiversity Strategies and Action Plans shows only 30 percent included actions to conserve and use agrobiodiversity. Considering 38% of land is now farmed and the importance of agrobiodiversity for wild biodiversity and vice versa, the absence of actions to increase and mainstream agrobiodiversity should be a big concern not only for the sustainability of food systems but also for the conservation of wild biodiversity. A central premise of the ABD Index is that loss of agrobiodiversity creates material risks for food and agricultural supply chains, for example shocks and long-term sustainability of sources of supply in the face of climate change and changing pest and disease loads and reputational risks with investors and consumers prioritizing healthy food from sustainable food systems. By detecting material agrobiodiversity-related risks and opportunities, the ABD Index is intended to motivate food and agriculture companies and public and private investors to mainstream agrobiodiversity in supply chains (e.g. diversified sourcing strategies) and financial instruments (e.g. positive or negative screening). The basic structure includes three pillars: 1) diets and markets seeks to capture to what extent and how food and agriculture companies, countries and projects contribute to ensure food biodiversity for healthy diets. 2) production systems seeks to capture to what extent and how food and agriculture companies, countries and projects contribute to agricultural diversity for sustainable production. 3) genetic resources seeks to capture to what extent and how food and agriculture companies, countries and projects contribute to diverse genetic resources for current and future/equal options towards adaptation and innovation. The ABD Index relies on three levels of measurement: 1) commitment: scores the level of commitment to agrobiodiversity as expressed in publicly available documents; 2) actions: scores the performance for a list of agrobiodiversity supporting practices; 3) status: the actual measurement of agrobiodiversity in terms of species, varieties, functions and specific landscapes features. To support this, we are building up a spatial layer of georeferenced data points. The evidence base and datasets that feed into this are large and growing. **Metrics** 33 indicators feed into the ABD index - each calculated based on a number of actual measurements, of which there are 76 in total at the moment. Some are more accessible for the current version. The colour coding indicates current availability of data. Category Indicators Markets & Production Genetic Source Consumption resources Status Species diversity Public and private Varietal diversity* 7 indicators datasets, evidence-Functional diversity base modelling, remote Underutilized/local species sensina. Soil biodiversity crowdsourcing Pollinator biodiversity Landscape complexity Reports, Global Actions Consumption practices supporting ABD 5 indicators datasets Avoided negative impacts Favor the maintenance of ABD Enhance use of ABD Supporting GR conservation Commitment Level of commitment based on 21 Policies, strategies, 21 indicators indicators declarations, etc. *Not available from secondary Largely available and feasible Partially available and needs work sources/prototype stage Outcomes Primarily based on real data. Soil and pollinator diversity are based on extrapolation based on through meta-analysis using the PREDICTS model. modelling















THE AGROBIODIVERSITY INDEX (ABD index) FICHE NR 6 Partnering with a commercial agricultural lender to develop an ABD Index score for their loan portfolio could stimulate ABD Index scoring and reporting by multiple agricultural lenders, eventually enabling sector-wide ABD Index reporting. Advising a company on its internal ABD Index monitoring system and agrobiodiversity enhancement in its value chain strategy could, if successful, encourage similar endeavors by multiple companies in a sector or region. If material benefits are observed, this could lead to development of an ABD Indexrelated data platform by several frontrunner companies in a sector or region. Successful issuance of an ABD Index-scored green bond (e.g. in partnership with a development finance institution) could lead to additional ABDI-scored green bonds within a region or sector and further to development of an investment fund of ABDI-scored bonds. Development of an ABD Index country score (e.g. in partnership with the Global Environment Facility) could provide the foundation for developing ABD Index scores for multiple countries (e.g. GEF clients) in a region, eventually enabling countries with higher or improved ABDI score to gain preferential access to sustainability finance (e.g. GEF biodiversity financing). Product line: Uptake at scale and e: Each color represents an example product derived from the ABD Index blications developed and scaled through three phases Minimum viable product: Early design and testing Required Three types of data feed into the ABD Index: data Publicly available reports, strategies, policies, product information Privately shared information on geospatial location of activities Global geospatial datasets and other global datasets. Granularity Granularity varies for specific indicators that feed into the Index; but the Index as a level whole is calculated at project, company or country level. User The ABD Index brings a wide variety of data together in order to make the friendliness information that those data provide, more actionable and usable by companies, countries and investors. The online portal presents the data in simple graphics and also allows deeper dives into details where of interest. Strengths The focus on **agro**biodiversity, being very complementary to indices that focus on recognised wild biodiversity and contributing also directly to better identify and manage by initiators **operational risks** – related to low agrobiodiversity, alongside reputational risk. The food system approach of the ABD Index, i.e. including three connected pillars: diversity in markets and consumption for healthy diets, diversity in production systems for sustainable agriculture, and diversity in genetic resources for current and future options.







FICHE NR 6	THE AGROBIODIVERSITY INDEX (ABD index)
	 The architecture that includes commitments, actions and status, allowing identification of where change is needed/ possible. The relatively low need for data input since most information is coming from publicly available documents and datasets.
Weaknesse s recognised by initiators	 Specific data gaps, particularly varietal diversity – identified as a critical aspect to make progress on. The limited quality or resolution of some secondary databases – identified as a critical aspect to make progress on. The aggregated ABD Index can be presented and communicated in a relatively simple way but the underlying measurements and indicators are quite complex.







8. The Biodiversity Footprint Calculator (PLANSUP)

FICHE NR 7	BIODIVERSITY FOOTPRINT CALCULATOR							
Date of assessment	 First description by EU B@B platform: August 2018 Complete description by Plansup: 4 November 2018 							
Actors	<u>Lead</u> : This calculator web tool was made by Plansup in collaboration with Saxion. <u>Other</u> : The Calculator is based on the Biodiversity Footprint Method which is developed by Plansup in collaboration with Wageningen Environmental Research (Alterra) and Netherlands Environmental Assessment Agency (PBL)							
Process and current position	The Calculator is a limited version of the full Biodiversity Footprint Method and is operational. The method is based on international recognized scientific research and has been implemented in several company cases and one agricultural sector. The development of the methodology has been commissioned by the Dutch Platform on Biodiversity Ecosystems and Economy (Platform BEE).							
	The research of the underlying methodology and the development of the calculator was carried out with public money and therefore it is fully accessible and free of charge. It is expected that the tool will be improved in the future on the basis of additional research and feedback from businesses.							
Key features of methodology	The method can be used to assess both current and future (or alternative) biodiversity footprint of a company's product at the landscape level. With the tool companies assess							
Matrice	The full Biodiversity Footprint Method (BFM) (see http://www.plansup.nl/expertise/biodiversity-footprint/) calculates the impact of the three most important terrestrial pressure types land use, greenhouse gas emissions and water use, and the impact of one of the most important aquatic pressure types: emission of nitrogen and phosphorus in (inland) water. The impact is calculated for all parts of the product chain. The Biodiversity Calculator is a limited version of the BFM and designed as a simple web tool that can be used by any company that would like to start with a quantification of their main impact on terrestrial biodiversity and that would like to test the effectiveness that different measures have on biodiversity, The calculator includes the impact of land use and greenhouse gas emissions and calculates the impact for the three most important parts of the chain: raw materials, production process and transport. The impact of onsite water use and chemical pollutants is included in the impact calculation of land use (part of land use management). With the help of this tool, companies have an opportunity to calculate their biodiversity footprint online. The tool uses dose response relationships from the GLOBIO methodology. With the calculator it is possible to calculate the biodiversity footprint for multiple scenarios. The tool enables companies to prioritize impact reducing measures and helps to determine which of the measures are most cost effective. Results are generated real time in both graphical and tabular format and can be downloaded. The tool itself and more information can be found on http://www.plansup.nl/biodiversity-footprint-calculator/ Important aspects of the methodology are its location specificity and use of real company data. The Biodiversity Footprint Method and Calculator can be used at product level for individual companies.							
Metrics	The indicator combines the area of the impact with the loss of biodiversity in the impacted area, expressed in MSA.ha. The MSA indicator is used in the GLOBIO methodology and can be expressed as an indicator for 'naturalness'. An MSA of 1 indicates no change (primary/original ecosystems) and 0 for completely degraded ecosystems (e.g. asphalt).							







FICHE NR 7	BIODIVERSITY FOOTPRINT CALCULATOR						
	For the biodiversity footprint the MSA is multiplied with the extent of the impacted area. The larger the footprint, the higher the impact of the product (/company) on nature.						
Outcomes based on modelling or real data?	The methodology uses primary company data as input. Only if a company does not have all data available, information can be extracted from existing relevant LCA studies and literature. The output is generated based on modelled dose response data.						
Drivers of biodiversity loss	Full Biodiversity Footprint Method: Land use (including on site water use and chemical pollutants) and Greenhouse Gas emissions, Water use, Nitrogen and Phosphorus emissions in water. Biodiversity Footprint Calculator: Land use (including on site water use and chemical pollutants) and Greenhouse Gas emissions						
Application	The method can be used for assessments at product, company and sector level. The calculator offers a calculation of the biodiversity impact of a products supply chain (raw materials), production process and transport. The full method also includes the other parts of the chain. The impact of multiple scenarios can be calculated and used to compare the effectiveness of potential or implemented measures but also for existing and new operations.						
Required data	Calculator: Company and chain related data in relation to land use and greenhouse emission (per pressure, location)						
Granularity level	Product, company or sector						
User friendliness	Calculator is made as user friendly as possible by only focussing on the main pressures and main parts of the supply chain. The data can be filled in by non-experts. All calculations are transparent and explained in the tool. The tool also provides instructions on what to do in case data are missing. Once the different types of land use and GHG emitters are identified for each of the three parts of the chain the user can start with the tool. In case land use can be described in general terms including extent (ha), and emitter types area known (type of energy use, transport type) the user can implement the calculator within one day. Preferably company emission figures are known for the relevant production processes. If not, existing general emission data will have to be used from similar processes (e.g. from LCA's)						
Strengths recognised by initiators	 Link between pressures and impacts (dose response relations) internationally used and scientifically approved (GLOBIO) Location specific Covers the most important drivers for biodiversity loss (Land use and GHG cover more than 80% of biodiversity loss; (Based on LCA research Netherlands in Report CE Delft Bergsma et al 2014 'The impact on biodiversity by Dutch companies') Biodiversity input data (MSA, pressure – impact relationships) based on extensive meta-analysis which continuously allows for adding new studies Allows for introducing weight factors differentiating ecosystem condition based on protection regime, protected species, etc. Includes economic allocation corrections for multiple use 						
Weaknesses recognised by initiators	 No impact on marine biodiversity. Impact is on general biodiversity and not species specific Impact of water use (extraction) on biodiversity neighbouring water sensitive nature areas only for Dutch situation Impact on site toxic emissions and water use is indirectly included as part of the impact of land use and cannot be assessed separately. Calculator does not (yet) calculate impact of water use (on neighbouring nature areas) and N & P emission in water. The latter is expected to be included in a follow up version of the calculator. 						







9. The LIFE Impact Index + Positive Scoring (by LIFE INSTITUTE)

FICHE NR 8	LIFE Impact Index + Positive Scoring						
Date of assessment	• First version 20/08/2018						
	Update on 2/11/2018						
Actors	Lead: LIFE Institute LIFE Institute is a non-profit organization headquartered in Brazil who operates internationally to the development and implementation of LIFE Methodology and its environmental managements tools.						
	LIFE Institute is also responsible for the operationalization of LIFE Certification System and the accreditation of independent Certifying Bodies.						
	Other: LIFE Permanent Technical Committee Committee composed by representatives of various organizations from different sectors: business, civil society and academia, as well as environmental government agencies.						
	Technical Committee members voluntarily contribute to the development and promotion of LIFE initiative.						
	A collaborative project is being established to adapt LIFE Methodology to Europe. The European actors for LIFE implementation process will be defined by the first half of 2019.						
Process and current position	LIFE Methodology is an international tool developed by LIFE Institute, which guides and recognizes businesses organizations that promote effective Natural Capital conservation actions contributing to the maintenance of Biodiversity and ES.						
	LIFE Methodology helps organizations in identifying their impacts and designing a strategic plan to reduce, mitigate and compensate them, including a specific approach to reduce impacts in the supply chain. LIFE's main objective is to disseminate and scale-up Natural Capital conservation and to contribute to make production practices more sustainable.						
	LIFE Methodology was designed as a practical and pragmatic tool to be applied in companies of any size and sector, adaptable to every country or region.						
	LIFE Methodology can be used both as an Environmental Management System (LIFE Key Software) and as a third-party Certification Scheme .						
	The initiative is currently operational in Brazil and Paraguay, with an expansion plan in execution in Latin America and Europe. Some 28 companies/ organizations are LIFE users and/ or have carried out technical projects to improve their environmental performance based on LIFE Methodology. In Brazil, 5 companies are already LIFE Certified, as well as 1 company in Paraguay.						
	LIFE is aligned with the Convention on Biological Diversity (CBD) and its Aichi targets as well as other international organizations and initiatives.						
	More information about LIFE initiative and all technical documents can be found under https://institutolife.org/en						
Key features of methodolog y	The Methodology is composed of complementary quantitative and qualitative approaches for a more comprehensive and complete impact/performance analysis; together with a more effective design of an action plan focused on concrete results.						
	A LIFE company is mainly committed to:						








Metrics

The quantitative approach is based on the calculation of an **Impact Index on Natural Capital** and consequent definition of a **Minimum Score in positive Actions**. consistent with the calculated impact.

These calculations are automatically performed by LIFE Software (LIFE Key) once the requested data are provided.

This analysis is complemented with the use of LIFE Matrix of Impacts on Biodiversity and ES, resulting from the company's operations. This module helps the identification/ analysis of business impacts as well as the dependencies, risks and opportunities related to them.



As part of the process of adapting the tool to Europe, the metrics will be reviewed and some adjustments will be possible. A broader scope of positive actions and additional technical developments are also foreseen to meet the needs of European companies and initiatives.

Natural Capital Impact Index

The calculation of LIFE Natural Capital Impact Index is based on selected environmental aspects that can be measured (or estimated) by any type of company: **Waste generation**, **GHG emission**, **Water consumption**, **Energy use** and the **Area occupied** by the operations. The impact of each one of these aspects is calculated in terms of **quantity and severity** (see picture below).







FICHE NR 8 LIFE Impact Index + Positive Scoring

A company fulfills the requested data and the system calculates **the resulting impact index for each aspect, as well as a final average impact index**. To this end, LIFE software must be customized by country or region and fed with all Natural Capital information required to the calculations. Data is obtained from official documents and bodies/ agencies (or internationally recognized organizations). Some examples of required information (by country or region) are: total GHG emissions, total waste generated, energy and water consumed per year, water availability per region, hierarchy of fragile and threatened ecoregions.

Index calculations are adjusted considering the maximum values known (per company) in the country or region.

Aspects	Waste	Greenhouse Gas	Water	Energy	Area
Quantity	Total annual amount generated (t/year)	Total annual emissions	Annual volume used (m³/year)	Total annual amount consumed (tep/year)	Total area occupied by operations (ha)
Severity	Hazardousness Destination	amount (tCO ₂ e/year)	Balance between water availability and demand	Impact of the consumed energy sources	Fragility of the ecoregion

Specificities of the Impact Index calculation

Some recognized tools are used for the estimation of few aspects that companies cannot directly measure (specific scientific studies on estimations can also be used):

- Water Footprint Network
- GHG Protocol and GHG Protocol Agricultural Guidance
- Regional evapotranspiration of forest plantations estimation (water withdrawn from the natural system)







FICHE NR 8 LIFE Impact Index + Positive Scoring In addition, the MSA (Mean Species Abundance) GLOBIO index is applied to better analyse biodiversity in the area occupied by the company. Estimated Evapotranspiration - IPEF MSA Water Footprint of Crops- WFN UNEP / WCMC Greenhouse Gas Aspects UNEP WCMC Blue Water Total area Green Water occupied by Quantity Annual Total annual operations (ha) volume used emissions (m³/year) amount (tCO2 e/vear) Balance between water Fragility of the Severity availability and ecoregion demand GHG GHG for Agriculture Minimum Performance in Natural Capital Conservation Actions LIFE tool calculates a Minimum Performance in Natural Capital Conservation for the voluntary compensation of residual impacts. Calculation of the minimum performance scoring considers the above-mentioned Impact Index. Scoring points obtained by means of conservation actions are defined according to an established list of actions/ projects, hierarchically ranked according to national or regional priorities for conservation. The scoring hierarchy system focuses on projects providing more concrete and effective results, considering technical qualifiers, indicators and the duration of the action. The results indicators define minimal monitoring priorities that capture changes in Biodiversity and Ecosystems Composition, Structure and Function. In the context of adapting the methodology for Europe, positive actions assessed by LIFE will be reviewed to ensure the inclusion of suitable Natural Capital conservation actions to fit European needs, priorities and programs. The calculation of a minimum positive performance is a parameter for: Analysing the adequacy of positive actions already implemented by the company and/ or Establishing an Action Plan on and /or Meet a LIFE Certification requirement Outcomes Hybrid. In most cases data are real and updated annually. Exceptionally, when real data based on are not available, estimations can be used and should be replaced over time by actual modelling or monitoring data. real data? **Drivers** of Drivers of biodiversity loss include: land conversion, fragmentation, climate change, biodiversity pollution, large environmental disturbances. Drivers have been added to the methodology loss as a result of LIFE continuous improvement. **Application** LIFE Methodology can be used both as an Environmental Management System and/ or as a Certification Scheme.

It applies to companies and organizations of all sizes and sectors, considering a **site** assessment approach taking into consideration the scope of the company's management







FICHE NR 8 LIFE Impact Index + Positive Scoring responsibility as well as the place(s) where conservation actions are implemented. The methodology also contains specific guidelines for reducing supply chain impacts. The site to be analysed is defined by the company. LIFE can be used to assess an entire group/ holding, a specific brand/ company department or a business unit. LIFE Methodology/Software must be fed with national data prior to its use. LIFE Management System Using LIFE Key software, the company can access the complete methodology or specific modules that it deems relevant. In this case, the company is considered as a LIFE user. LIFE Certification A company that complies with LIFE indicators and achieves the Minimum Performance in Conservation Actions may request a third-party audit to obtain LIFE Certification. Certification Bodies must be accredited by LIFE Institute. All documents are available on LIFE website. Main technical documents are: LIFE Standards (Qualitative assessment: environmental management indicators) LIFE Technical Guide 01 (Calculation of the impact index) LIFE Technical Guide 02 (Scoring system for conservation actions) Note: The evaluation of impacts on biodiversity and ES module is not yet available online Required data Company data: Sector of activity, business structure, unit (site) selected for the methodology application, identification/ location information, annual income **Environmental Information:** • Waste - type of waste, destination and annual quantity generated · Gas - gases emitted, GHG Protocol scope (1,2,3), annual quantity emitted • Water - type of water source, annual amount consumed • Energy - energy source, annual quantity consumed • Area - ecoregion, type of land occupation (agriculture, built area, etc.), area size Environmental projects and conservation actions carried on Type of action (according to LIFE Methodology) Ecoregion Purpose of the action Description of the action · Initial and final date · LIFE qualifiers and indicators attended Impacts analysis - Biodiversity and ES · Processes, activities and their resulting impacts, affected ES, dependencies, risks and opportunities (the completion of the matrix is guided by options lists and orientations provided by the tool) Granularity Business unit scaled to the country or region level User Is it easy to use by non-experts or does it require specialist knowledge? friendliness LIFE Key software Is easy to use by non-experts and does not require specialist knowledge. A quick training is recommended for a better understanding of the tool. However, a specialist can help to refine and detail the analysis and the elaboration of the action plan.







FICHE NR 8	
	Software accessible via the Internet. Navigation based on Windows and Office templates.
	What are the required time efforts? Efforts depend on the level of detail sought, especially in relation to conservation projects and action plan. The first full and detailed assessment should take a few months and annual updates are much quicker.
	The different modules of the methodology can be used separately for specific evaluations.
	Does it require a similar approach as other environmental issues or is it very specific? LIFE is not a sectoral specific methodology.
	Some tools and scientific studies are used as part of LIFE calculations: Water Footprint Network; GHG Protocol and GHG Protocol Agricultural Guidance; regional evapotranspiration of forest plantations studies. MSA (Mean Species Abundance) GLOBIO index is applied to better analyse biodiversity in the area occupied by the company.
Strengths recognised by initiators	 Focused on Biodiversity and ES conservation Pragmatic tool applicable to organizations of any size and sector Complementarity between qualitative and quantitative approaches Designing of strategic conservation plan scenarios Measures, evaluates, monitors and compares business positive and negative impacts Enables comparability between companies, business units and sectors Issues executive and detailed reports for clear and transparent results communication to different stakeholders Possibility of recognition by third-party Certification Strategic orientation to assure investment return in conservation by focusing on concrete actions
Weaknesses recognised by initiators	 Need for prior adaptation by country or region Visibility of returns on investment (business as usual perspective) Scaling-up pace Still to be developed a financial valuation module (negative/ positive impacts and dependencies on ES)







10. Bioscope (by Platform BEE, Dutch Ministry)

FICHE NR 9	BIOSCOPE
Date of assessment	30 August 2018 (assessment by EUB@B Platform, based on information on internet)
Actors	<u>Lead</u> : BioScope has been developed by PRé Sustainability, Arcadis and CODE, commissioned by Platform BEE (Biodiversity, Ecosystems and Economy); a collaboration between IUCN NL and VNO-NCW financed by the Dutch ministry of economic affairs.
Process and current position	Operational
Key features of methodology	BioScope provides businesses with a simple and fast indication of the most important impacts on biodiversity arising from their supply chain .
	The results brought by BioScope are aimed at helping you to formulate meaningful actions to further assess and reduce the impact of your business on biodiversity. It not only indicates the potential impact of the commodity you purchase, but also of the upstream supply chain of these commodities . BioScope makes use of Exiobase v2.2 enabling you to select commodities and resources purchased from 170 sectors in 43 countries (27 EU countries and all large economies outside the EU), covering the largest part of global economic activities. The resulting impacts on biodiversity are calculated with the ReCiPe method, which was specially adapted for BioScope.
	More info can be found in the Methodological Guidance, see https://www.bioscope.info/uploads/bioscope.info/bee_downloads/9/file/Methodology_Report_v1.compressed.pdf
Metrics	PDF.m2.yr (for land) and PDF.m3.yr (for water). PDF stands for Potentially Disappeared Fraction of species.
Outcomes based on modelling or real data?	Modelling
Drivers of biodiversity loss	Most important drivers for biodiversity loss are taken into account, including: climate change, terrestrial ecotoxicity, terrestrial acidification, land occupation, land transformation, water use/scarcity, marine ecotoxicity, fresh water eutrophication and fresh water ecotoxicity. Impact drivers not included in the quantitative assessment include the introduction of invasive species and disturbance.
Application	 Supply chain decisions. Examples of questions which can be answered are: Which of the commodities purchased by my business could be the largest cause of impact on biodiversity? What could the new purchasing strategy of my business mean for our impact on biodiversity? What commodity purchased by my business do we need to focus on if we want to make a meaningful contribution to conservation of biodiversity? In which regions are these impacts localized?
Required data	An inventory of commodities used in a given supply chain; this is done by specifying the expenditure per commodity for each stage. A biodiversity impact mode (ReCiPe), which will translate these regionalized economical activities to meaningful indicators that describe their influence with regard to each of the impact drivers
Granularity level	Parcel level
User friendliness	Very user friendly (see Quick Start Guide







FICHE NR 9	BIOSCOPE
	https://www.bioscope.info/uploads/bioscope.info/bee_downloads/8/file/User's_quick_start_v1.pdf
Strengths recognised by initiators Weaknesses recognised by initiators	Strengths and limitations of underlying Exiobase and ReCiPe models.







ANNEX 2: ANALYSIS OF ASSESSMENTS

The following tables list the elements of certain key features in a concise form that we consider vital for enhancing the transparency, understanding the approaches and for an initial comparative analysis. They may be the basis for a more thorough analysis at a later stage.







Annex 2 Table 1: Key features of assessed initiatives on biodiversity metrics for business

	State of art	Sectors	Metric	Key elements of methodology	Real data	Which data
	Full	All sectors,	Mean species	Two-step process:	or modelling Hybrid.	Economic
1. Global Biodiversity Score	development by 2020. Operational relevance being tested.	including FI	abundance (MSA) and its surface area equivalent, i.e., km² MSA	 Quantitative assessment of pressures caused by specific economic activities on biodiversity; value chain analysis based on Exiobase matrix-based input-output model and direct data on pressures when available. Estimation of impacts of these pressures on ecosystems; this relies on the GLOBIO model which is based on pressure-impact relationships. GLOBIO includes land conversion, fragmentation, encroachment, atmospheric N deposition (eutrophication) and climate change for terrestrial biodiversity, and wetlands conversion, local and network landuse in catchment of wetlands, hydrological disturbance of wetlands and rivers, land-use in catchment of rivers and eutrophication of lakes for aquatic biodiversity. Missing drivers will be added to GLOBIO through future developments 	Real data on pressures can be included. When very detailed ecological monitoring data are available, the mean species abundance might be directly calculated.	activity data: turnover by country and industry (of the company assessed or of the company a financial institution invested in); Pressure data: - Carbon emissions on scope 1, 2 and 3 - Land use changes (ideally using a 13 habitat types nomenclature including different use intensity for forests, grasslands, agriculture, etc.) Comprehensive biodiversity direct data
2. Biodiversity Impact Metric	Pilot testing of beta version.	Particularly for MNCs that source	The Biodiversity Impact Metric	The methodology is focused on measuring the impacts of raw material production, or land	Hybrid.	Company land area Proportion of biodiversity lost





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
	Extending the possibilities in the future.	hundreds of different raw materials from across the planet	is quantifiable measure that can be used to assess and track the impact of a company's land use activities on biodiversity in a given area.	use, in global supply chains. The basic framework for the metric is as follows: Biodiversity impact = land area x quantity impacted x quality impacted Area (ha) of land needed to produce commodity (Company data on amount of commodity and source location Data on country level yield The various components in the metric are assessed at the eco region level. Metric results are calculated at the scale of Country Ecoregion Component and then aggregated to produce a national score for different commodities.		(MSA coefficients for different land uses and management intensities) The importance of a location for biodiversity, based on range rarity scores (underpinned by IUCN Red List data)
3. Biodiversity Indicators for Extractive Companies	Under development	Extractive industries (O&G, mining)	Set of site level indicators, allowing aggregation at corporate level (to be developed). It was decided that a single metric may not provide the necessary resolution, flexibility or information to meet the indicator needs of the	A three stages process is suggested for indicator development: 1. First stage: screening of the company's portfolio of operations to identify sites with potentially high biodiversity sensitivity, based on globally and locally available data sets combined with site validation to identify sensitive sites for prioritization of reporting effort; 2. Second stage: development of site level biodiversity indicators using the state-pressure-response (SPR) framework (a widely accepted organising framework for biodiversity management and monitoring), informed by the stage above and based on site level data and documentation for	It is intended that the approach uses real data, but modelled/ estimated data may also be used. This will be explored within the piloting process	Red List species range rarity layer - 1x1km resolution (IUCN). Available through IBAT as of January 2019 Global Critical Habitat screening layer (UNEP-WCMC). Available through IBAT as of January 2019 Polygon-based - World Database on Protected Areas (IUCN and UNEP-WCMC). Available through IBAT





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
			extractive sector.	high sensitivity sites collected as part of the environmental impact assessment; and		
				Third stage: aggregation of scores for SPR at site level up to business unit, division, and corporate level to provide insight into performance on the ground.		
	Pilot phase. Further improvements in 2018/2019	All industries	Potential disappeared fraction of species within a year, or PDF*yr (frequently used in LCA methodologies)	The main objective of the project is to improve the biodiversity performance of a product by identifying biodiversity hotspots that can be improved and support eco-design approaches. The LCA framework is used to calculate the relative differences between the variants of a product. In the long term, it will also allow to compare different products or different sectors at larger scales. The methodology is based on 3 modules	Hybrid approach, based on modelling and real data	LCI Data (EcoInvent, Agribalyse,) Biodiversity database (IUCN, IBAT, Predicts,) Data provided by company on biodiversity measures
4. Product Biodiversity Footprint				 Module 1 computes lifecycle impact assessment, with spatial differentiation for the main impact categories evaluated. Characterization factors used for computation are the ones currently available in published LCA methodologies (LC Impact). This first module enables the user to visualize the hotspots of the product footprint both geographically and along the whole value chain. Module 2 treats specific information regarding the practices and the local context, that would enable to adjust 		





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
				impact computations based on information entered by the users. The development of criteria and rules to quantify the changes in impact computations are defined per type of land use (e.g. arable crops, perennial crops, grassland, mining, forest, urban) and/or per sector (livestock, transport, electricity, construction). Results of module 2, additionally to the results of the Module 1, enable the user to visualize and quantify the benefits of a chosen practice/location, and compare various scenarios for a given product. - Module 3 assesses qualitatively 2 aspects that are not part of any LCA model, namely 'invasive species' and 'species management', this last one encompassing 'overexploitation', but going beyond as it also includes positive actions (e.g. installation of pollinators, use of various breeds, follow up of endangered species).		
5. ASN Biodiversity Footprint approach	Developed and being applied for 2 years now	FI, covering investments in all sectors	PDF.m2.yr (for land) and PDF.m3.yr (for water). PDF stands for Potentially	The quantitative methodology consists of 3 steps: The <u>first step</u> is to create an overview of the economic activities the FI invests in. This step includes:	Real data from Exiobase to calculate the environmental footprint on a	Data on economic activities of companies invested in. Exiobase data; the use of other data-





State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
		Disappeared Fraction of species. The PDF.m2.yr and PDF.m3.yr can be added up to a PDF.yr or species/yr score (using the average species density on land and in water).	 A 'definition' of the activities of a company: what is the company producing (in what sectors is the company active?) and where does production take place? Decisions on the scope of the assessment (supply chains included or not); A selection of the investments included in the assessment (all major investments) In the second step the environmental impact of the economic activities of the companies invested in is assessed. The environmental data in the 'Exiobase' input-output database²⁹ is used to assess what land use, water use, emissions, etc. is linked to the economic activities of the companies. Exiobase takes into account world-wide trade flows between countries and between sectors. In the third step, the ReCiPe methodology is used to calculate the environmental footprint on a midpoint level (e.g. climate change resulting from CO2 emissions) and to calculate the resulting impact on ecosystem quality or biodiversity (endpoint level). This latter step is based on science based 'doseresponse' relations (e.g. the effect of a 1 degree temperature rise on biodiversity). This results in an impact on terrestrial biodiversity and an impact on aquatic biodiversity. The result is then used to calculate the biodiversity footprint in m2 per Euro invested (for each 	midpoint level and dose-response modelling to calculate the biodiversity footprint of economic activities / companies. No primary data (actual data from the companies invested in) are used in this step, only secondary data (sector averages from Exiobase).	sources (like Trucost data) is being explored. Information on biodiversity impact drivers in different sectors to allow for a qualitative analysis guiding interpretation and use of the footprint results

²⁹ https://www.exiobase.eu/





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
				investment category) and the total footprint in m2 for all investments. A gualitative analysis is used to guide the interpretation and the use of the footprint results, looking at (among others) the limitations of the footprinting methodology and the potential influence of the footprint results on investment decisions.		
6. Agrobiodiversity Index (ABD)	prototype Version 1. Elaboration of case studies and continuous fine-tuning	Agro- industry	ABD Index based on 33 indicators. Although only a few indicators relate to wild biodiversity, the index could be considered as a proxyindicator for wild biodiversity.	The focus is on agricultural biodiversity at the genetic, species and landscape levels, i.e. the variety and variability of animals, plants and micro-organisms that are used directly or indirectly for food and agriculture (FAO definition). This makes the ABD index very complementary to other metrics or indices that focus primarily on wild biodiversity. The basic structure includes three pillars: 1) diets and markets seeks to capture to what extent and how food and agriculture companies, countries and projects contribute to ensure food biodiversity for healthy diets. 2) production systems seek to capture to what extent and how food and agriculture companies, countries and projects contribute to agricultural diversity for sustainable production. 3) genetic resources seek to capture to what extent and how food and agriculture companies, countries and projects contribute to diverse genetic resources for current and future/equal options towards adaptation and innovation. The ABD Index relies on three levels of measurement: 1) commitment: scores the level of commitment to agrobiodiversity as	Primarily based on real data. Soil and pollinator diversity are based on extrapolation through meta- analysis using the PREDICTS model.	Three types of data feed into the ABD Index: - Publicly available reports, strategies, policies, product information - Privately shared information on geospatial location of activities - Global geospatial datasets and other global datasets





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
				expressed in publicly available documents; 2) actions: scores the performance for a list of agrobiodiversity supporting practices; 3) status: the actual measurement of agrobiodiversity in terms of species, varieties, functions and specific landscapes features.		
7. Biodiversity Footprint Calculator	Operational	All sectors	MSA.ha (Mean Species Abundance)	A free calculation tool to assess both current and future biodiversity footprint of a company's product at the landscape level. With the tool companies can test the effectiveness of presumed biodiversity friendly measures. With the help of this tool, companies have an opportunity to calculate their biodiversity footprint online. The tool is based on dose response relationships of two pressure factors from the GLOBIO methodology and therefore it gives only an indication of the generic impact on biodiversity. This tool calculates the biodiversity impact of a company's supply chain, production process and transport that can be related to one or more products. With the calculator it is possible to calculate the biodiversity footprint for multiple scenarios. It is recommended to use the first scenario to describe the current situation and to use subsequent scenarios to describe all kinds of biodiversity friendly measures that are expected to change the footprint in the near future.	Modelling	Company data (location, pressures GHG and land use) GLOBIO data
8. Impact Index + Positive Scoring	Operational	All sectors	MSA (Mean Species Abundance), as part of a	LIFE Methodology helps organizations in identifying their NC impacts and designing a strategic plan to reduce, mitigate and compensate them, including a specific	Hybrid. In most cases data are real and updated	Company data Environmental information:





Si	state of art Sec		Key elements of methodology	Real data or modelling	Which data
		ider NC npact Index	approach to reduce impacts in the supply chain. LIFE Methodology can be used both as an Environmental Management System (LIFE Key Software) and as a third-party Certification Scheme. Quantitative approach is based on the calculation of an Impact Index on Natural Capital and consequent definition of a Minimum Score in positive Conservation Actions. consistent with the calculated impact. Calculations are automatically performed by LIFE Software (LIFE Key) once the requested data are provided. This analysis is complemented with the use of LIFE Matrix of Impacts on Biodiversity and ES, resulting from the company's operations (based on GLOBIO pressure – impact relationships). LIFE tool calculates a Minimum Performance in Conservation Actions for the voluntary compensation of residual impacts. Calculation of the minimum performance scoring considers the outcomes of the NC Impact Index. Scoring points obtained by means of conservation actions are defined according to an established list of actions/ projects, hierarchically ranked according to national priorities for conservation. The scoring hierarchy system focuses on projects providing more concrete and effective results, considering technical qualifiers, indicators and the duration of the action. The	annually. Exceptionally, when real data are not available, estimations can be used and should be replaced over time by actual monitoring data	Data on pressures and area (ecoregion, type of land occupation (e.g. agriculture, built area, etc.), area size Environmental projects and conservation actions Impacts analysis – Biodiversity and ES: Processes, activities and their resulting impacts, affected ES, dependencies, risks and opportunities (the completion of the matrix is guided by options lists and orientations provided by the tool)





	State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
				results indicators define minimal monitoring priorities that capture changes in Biodiversity and Ecosystems Composition, Structure and Function. LIFE Methodology/Software must be fed with national data prior to its use		
9. Bioscope	Operational	All sectors	PDF.m2.yr (for land) and PDF.m3.yr (for water). PDF stands for Potentially Disappeared Fraction of species.	BioScope provides businesses with a simple and fast indication of the most important impacts on biodiversity arising from their supply chain. The results brought by BioScope are aimed at helping you to formulate meaningful actions to further assess and reduce the impact of your business on biodiversity. It not only indicates the potential impact of the commodity you purchase, but also of the upstream supply chain of these commodities. BioScope makes use of Exiobase v2.2 enabling you to select commodities and resources purchased from 170 sectors in 43 countries (27 EU countries and all large economies outside the EU), covering the largest part of global economic activities. The resulting impacts on biodiversity are calculated with the ReCiPe method, which was specially adapted for BioScope.	Modelling	An inventory of commodities used in a given supply chain; this is done by specifying the expenditure per commodity for each stage. A biodiversity impact mode (ReCiPe), which will translate these regionalized economical activities to meaningful indicators that describe their influence with regard to each of the impact drivers
10. Biodiversity Return on Investment Metric (BRIM)	Under development. First road- testing.	FI + All sectors	BRIM Ex-ante ROI for a species, and/or for a site Composed of 1/% of total population at	BRIM measures the change in risk of species extinction attributable to investment Investment can change the scale and impact of the processes that cause species to be at risk of extinction- deforestation, over-exploitation	Real data and coefficients	Species range maps or maps of Extent of Suitable Habitat from the IUCN Red List of Species





State of art	Sectors	Metric	Key elements of methodology	Real data or modelling	Which data
		site, 2/ Red List category weighting, 3/ relative contribution of each pressure (P x w x R)	Based on the IUCN Red List of Threatened Species Based on quantitative categories and criteria – not expert opinion Incorporates compilation of data on range, habitats, threats, etc – not just a list		Threat status and threats applying to species from the IUCN Red List of Species (complemented by information from the Key Biodiversity Areas dataset) Polygon of investment intervention is needed (site, protected area, country, landscape, land-use unit) Site- or country-based measures require bespoke assessments using GIS applications and integration of threat and land-use layers







Annex 2 - Table 2: Type of business applications for each biodiversity metrics tool

	nex 2 - Table 2: Type of	Product	Project	Site	Supply chain	Corporate	Finance
1.	Global Biodiversity Score		Project level assessments are robust only for very large- scale projects (impacting thousands of hectares) and should generally be limited to internal purposes			Corporate level assessments, including estimation of biodiversity performance along the whole value chain of a company. The GBS can also be used by nonfinancial companies themselves to assess a refined footprint if detailed information is available. The GBS could be used for No Net Loss approaches at the corporate level.	Calculating the footprint of a financial asset portfolio as a basis for portfolio investment decisions by finance sector, as long as it remains at the level of sectors or companies (when enough company-specific data is available). The GBS could be used for No Net Loss approaches at the financial institution-level.
2.	Biodiversity Impact Metric	Establishing an overall potential impact score from commodity sourcing, and comparing potential impacts of different			Flagging geographic sources of potential elevated impacts in a commodity supply chain		Comparing potential impacts of different companies sourcing the same commodity(ies) generally.







		1			
	commodities – within a company's supply chains or more				
3. Biodiversity Indicators for Extractive Companies		Assessing and monitoring progress of biodiversity risks. Comparing options. Applying a NNL approach	Assessing and monitoring progress of biodiversity risks Comparing options. Applying a NNL approach Communication and reporting	Assessing and monitoring progress of biodiversity risks Applying a NNL approach Communication and reporting	Comparing biodiversity performance of companies within the extractives sector
4. Product Biodiversity Footprint	Product evaluation (PBF evaluates the impact of products or services) and subsequent decision making on sourcing of material, change of production practices (e.g. farming) or eco- design. Product communication				
5. ASN Biodiversity Footprint approach					Calculating the footprint of a financial asset portfolio, and subsequent portfolio investment decisions by finance sector, as





					long as it remains at the level of
					sectors or companies.
					Development of investment
					criteria based on insights in the
					main impact drivers of different
					asset classes and sectors.
					Use as a scoping step: to
					identify biodiversity impact
					hot spots on a portfolio level,
					enabling follow-up steps for a
					selection of investments.
					Use this footprint of the portfolio
					/balance sheet to identify
					hotspots and create a strategy
					to reach a No Net Loss
			Measuring supply	Measuring	Impact investing, i.e.
			chain	corporate	investments "made into food
			performance	performance.	and agriculture companies,
6 Agrabiadiyaraity					organizations, and funds with
6. Agrobiodiversity					the intention to generate a measurable, beneficial social or
Index (ABD)					environmental impact alongside
					a financial return.
					issuance of an ABD Index -
					scored green bond
	Calculation of				Soored green bond
	biodiversity				
	impact of a				
	company's supply				
7. Biodiversity	chain, production				
Footprint	process and				
Calculator	transport that can				
Calculator	be related to one				
	or more products.				
	Multiple scenarios				
	can be calculated				





	(application of				
	(application of measures).				
	measures).	Mossuring biodive	roity parformance		Investment decisions could be
8. Impact Index + Positive Scoring		Measuring biodiversity performance. It applies to companies and organizations of all sizes and sectors, considering a site assessment approach taking into consideration the scope of the company's management responsibility as well as the place(s) where conservation actions are implemented. The methodology also contains specific guidelines for reducing supply chain impacts. The site to be analysed is defined by the company. LIFE can be used to assess an entire group/ holding, a specific brand/ company department or a business unit.			Investment decisions could be guided by certifications, such as LIFE
9. Bioscope	Assessment of biodiversity performance of specific commodities in the supply chain	Supply chain decisions by corporates. Examples of questions which can be answered are: • Which of the commodities purchased by my business could be the largest cause of impact on biodiversity? • What could the new purchasing strategy of my business mean for our impact on biodiversity? • What commodity purchased by my business do we need to focus on if we want to make a meaningful contribution to conservation of biodiversity? In which regions are these impacts localized?			
10. Biodiversity Return on Investment Metric (BRIM)		Application at site scale requires some data gathering and complementary information	·	Assessment of potential for species extinction risk reduction across corporate footprint (mine sites, plantations,	Screening of potential biodiversity ROI across a portfolio (for instance, investments in a particular company value chain, for a site-based commodity)







		landscape	
		restoration)	







COLOPHON

CRITICAL ASSESSMENT OF BIODIVERSITY ACCOUNTING APPROACHES FOR BUSINESSES DISCUSSION PAPER FOR EU BUSINESS @ BIODIVERSITY PLATFORM

AUTHOR

Johan Lammerant

DATE

19 November 2018

ABOUT THE EU B@B PLATFORM

The EU B@B Platform is a forum for dialogue and policy interface to discuss the links between business and biodiversity at EU level. It was set up by the European Commission with the aim to work with and help businesses integrate natural capital and biodiversity considerations into business practices. The EU B@B Platform focuses its work on three thematic workstreams: Natural Capital Accounting, Innovation and Finance. ICF is supporting the European Commission in running the EU B@B Platform since 2013. Arcadis is leading the Natural Capital Workstream.

Arcadis Belgium nv/sa

Gaston Crommenlaan 8 bus 101 9050 Ghent Belgium 02 505 75 00

www.arcadis.com

